

# Growth, toxicity and oxidative stress of a cultured cyanobacterium (*Dolichospermum* sp.) under different CO<sub>2</sub>/pH and temperature conditions

Andreas Brutemark,<sup>1,2†</sup> Jonna Engström-Öst,<sup>1,2\*†</sup> Anu Vehmaa<sup>2</sup> and Elena Gorokhova<sup>3</sup>

<sup>1</sup>ARONIA Coastal Zone Research Team, Novia University of Applied Sciences & Åbo Akademi University, Ekenäs,

<sup>2</sup>Tvärminne Zoological Station, University of Helsinki, Hanko, Finland and <sup>3</sup>Department of Applied Environmental Science, Stockholm University, Stockholm, Sweden

## SUMMARY

Cyanobacteria blooms are a worldwide nuisance in fresh, brackish and marine waters. Changing environmental conditions due to upwelling, changed mixing conditions or climate change are likely to influence cyanobacteria growth and toxicity. In this study, the response of the toxic cyanobacterium *Dolichospermum* sp. to lowered pH (−0.4 units by adding CO<sub>2</sub>) and elevated temperature (+4°C) in an experimental set-up was examined. Growth rate, microcystin concentration and oxidative stress were measured. The growth rate and intracellular toxin concentration increased significantly as a response to temperature. When *Dolichospermum* was exposed to the combination of elevated temperature and high CO<sub>2</sub>/low pH, lipid peroxidation increased and antioxidant levels decreased. Microcystin concentrations were significantly correlated with growth rates. Our results show, although oxidative stress increases when exposed to a combination of high CO<sub>2</sub>/low pH and high temperature, that growth and toxicity increase at high temperature, suggesting that the cyanobacterium in general seems to be fairly tolerant to changes in pH and temperature. Further progress in identifying biological responses and predicting climate change consequences in estuaries experiencing cyanobacteria blooms requires a better understanding of the interplay between stressors such as pH and temperature.

Key words: cyanobacteria, growth rates, microcystin, oxidative status, pH levels, warming.

## INTRODUCTION

The ongoing global climate change caused by combustion of fossil fuels and the following carbon dioxide (CO<sub>2</sub>) emissions result in warming of the atmosphere and seawater. If emissions continue at the current speed, a 2–4.5°C rise is most likely to occur (IPCC 2013). In addition, up-welling events and changed mixing conditions can expose plankton to high CO<sub>2</sub>/low pH water (Feely *et al.* 2008; Barry *et al.* 2011). Estuaries and coastal areas in particular have a unique status in the future: they will be important natural laboratories concerning ecosystem changes caused by increased upwelling, fuelled by ocean acidification (Feely *et al.* 2010; Wootton & Pfister 2012). The present pH of the ocean surface has already declined by 0.1 pH units compared with pre-industrial levels, and is predicted to continue decreasing (Caldeira &

Wickett 2003; IPCC 2007). Therefore, the combined dynamics of CO<sub>2</sub>/pH and temperature is more relevant than changes in a single factor.

In this context, of immediate interest are the responses of nuisance species, such as toxic cyanobacteria that today pose environmental problems in many aquatic ecosystems, including the Baltic Sea (O'Neil *et al.* 2012). For example, warming and low pH (31°C/900 µatm pCO<sub>2</sub>) caused a significant increase in growth rates of the oceanic cyanobacterium *Trichodesmium* spp. (Levitan *et al.* 2010), consistent with results in previous studies (Barcelos e Ramos *et al.* 2007; Hutchins *et al.* 2007; Levitan *et al.* 2007). Similar responses have been measured for the toxic cyanobacterium *Nodularia spumigena* Mertens (Wannicke *et al.* 2012; Endres *et al.* 2013; Unger *et al.* 2013).

Although genetic and/or phenotypic adaptations to low pH alone, or in combination with other factors, have been investigated for most algal groups including chlorophytes (Collins & Bell 2004), coccolithophores (Feng *et al.* 2008), diatoms (Sun *et al.* 2011; Kremp *et al.* 2012), dinoflagellates (Fu *et al.* 2008, 2010), raphidophytes (Fu *et al.* 2007), and cyanobacteria (Fu *et al.* 2007; Levitan *et al.* 2010), little is known about oxidative stress responses in algae during warming and low pH. Reactive oxygen species (ROS) are produced as by-products during photosynthetic electron transport and metabolism in photosynthetic organisms (Apel & Hirt 2004). Primary producers, such as higher plants and algae, are constantly exposed to oxidative stress from changes in light and other environmental factors. If the cells exposed to oxidative stress are not able to handle it, a loss of protein function, membrane integrity and eventually cell death can result (Monaghan *et al.* 2009). To counteract or prevent damage by ROS, microalgae, including cyanobacteria, need an anti-oxidative mechanism. The knowledge of the ecological role of oxidative stress and antioxidants in cyanobacteria is limited (Latifi *et al.* 2009; Zilliges *et al.* 2011), and there is

\*To whom correspondence should be addressed.

Email: jonna.engstrom-ost@novia.fi

Communicating editor: Rei Narikawa

†Contributed equally.

Received 26 March 2014; accepted 4 September 2014.

a need to include these measurements when studying cyanobacteria responses to environmental factors.

The aim of the current study is to measure growth, toxicity and oxidative stress of a cultured cyanobacterium under different CO<sub>2</sub>/pH and temperature conditions. The study area of the Baltic Sea is a shallow semi-enclosed brackish water basin, with a restricted connection to the North Sea. Due to high anthropogenic nutrient load from the land, the Baltic Sea is at present highly eutrophicated (Fleming-Lehtinen *et al.* 2008). The pollution is manifested for instance as annual cyanobacteria mass-occurrences covering large areas consisting mainly of three taxa: *Aphanizomenon* sp., *Dolichospermum* spp. and *Nodularia spumigena*. *N. spumigena* is a well-known toxin producer that affects aquatic and terrestrial organisms, and may pose a health hazard for humans (Sellner 1997; Mazur-Marzec & Pliński 2009). *Aphanizomenon* sp. and *Dolichospermum* spp. are known to produce toxins in freshwaters (Sivonen & Jones 1999) and more recently, *Dolichospermum* has proven to be hepatotoxic in the Baltic Sea, producing microcystins (Halinen *et al.* 2007; Fewer *et al.* 2009). In this experimental study, we mimicked a cyanobacteria bloom and monitored how increased CO<sub>2</sub>/lowered pH and elevated temperature, as single factors and combined, affect growth, intracellular toxin concentrations and the oxidative status of the cyanobacterium *Dolichospermum* sp.

## MATERIALS AND METHODS

### Cultivation of the strain

The *Dolichospermum* sp. BIR257 strain (previously *Anabaena* sp., Wacklin *et al.* 2009) originates from the Gulf of Finland (Baltic Sea) and is a microcystin-producer (Halinen *et al.* 2007). The culture was grown as batch monocultures in Z8 medium (Kótai 1972) without nitrogen at 18°C in 13.7 μmol photons m<sup>-2</sup> s<sup>-1</sup> in a 16:8 h light: dark cycle. Even though the strain was not axenic, aseptic techniques were always used when handling the culture.

### Sampling and experimental set-up

We used a factorial (2 × 2) experimental design with four treatments: (i) Control: 17°C and normal pH, (ii) Low pH: low

pH (−0.4 units) and 17°C, (iii) High temp: 21°C and normal pH, and (iv) Low pH + high temp: low pH (−0.4 units) and 21°C (Table 1). Normal pH is defined as pH in filtered seawater at 17°C. Water from 5 m depth was collected from a nearby pelagic sea area, located at the south-western coast of Finland, in the vicinity of Tvärminne Zoological Station (59°50′ N, 23°15′ E, University of Helsinki). The treatments were prepared by first filtering the seawater (salinity 5.5) using Sartobran 300 filters (<0.2 μm; Sartorius Stedim Biotech GmbH, Göttingen, Germany). As filtration can shift the carbonate chemistry of the seawater, CO<sub>2</sub> bubbling should take place after filtration (Riebesell *et al.* 2010). Then the filtered seawater was pre-conditioned to reach the experimental temperatures (17 and 21°C). Subsequently, the treatments with low pH were supplemented with CO<sub>2</sub> gas until pH had decreased by −0.4 units (TUNZE pH controller 7070/2, Penzberg, Germany) according to predictions for 2100 (Caldeira & Wickett 2003; IPCC 2007). The low pH used corresponds to the approximate expected pH change given as a general prediction for up-welling water (Feely *et al.* 2008). Finally, filtered seawater was mixed with *Dolichospermum* sp. culture (target concentration: 100 μg C L<sup>-1</sup>). To avoid nutrient limitation during incubations, phosphate, nitrate (Table 2), vitamins and minerals were added in surplus, following the Guillard (1975) recipe. pH and temperature were measured from each bottle prior to closing the lid (9 replicates treatment<sup>-1</sup>). 1.2 L Duran clear airtight glass bottles with Melamin resin screw caps with polytetrafluoroethylene seals were used and attached to plankton wheels (1 rpm) in climate chambers with respective temperatures. Light intensity was 18 μmol photons m<sup>-2</sup> s<sup>-1</sup> at the top of the plankton wheels. Samples for dissolved nutrients, particulate organic carbon (POC), oxidative stress biomarkers and intracellular toxins were collected at the start. Three replicate samples per treatment (Control, high CO<sub>2</sub>/low pH, high temp, high CO<sub>2</sub>/low pH + high temp) for each variable were collected directly after mixing. These start samples were taken before the treatment water was filled into bottles. In order to evaluate potential differences between treatments, one set of replicates was sacrificed and samples were collected after 8 days of incubation (data not shown). Since no difference was detected, the experiment was prolonged. After 11 days of incubation clear differences in biomass were observed between treatments and the experiment was terminated. End samples were taken on

**Table 1.** Changes in pH and dissolved inorganic carbon (DIC) in the units with and without *Dolichospermum* sp. pH is reported as minimum to maximum range

	Without cyanobacteria		With cyanobacteria	
	Start	End	Start	End
<i>pH</i>				
Control	7.8–7.9	7.7–7.8	7.6–7.8	8.3–8.8
High CO <sub>2</sub> /low pH	7.5–7.5	7.5–7.5	7.4–7.5	8.0–8.7
High temp	7.8–7.9	7.8–7.8	7.8–7.9	9.0–9.3
High CO <sub>2</sub> /low pH + high temp	7.5–7.5	7.5–7.5	7.5–7.6	8.9–9.2
<i>DIC</i>				
Control	1.8 ± 0.0	1.8 ± 0.0	1.9 ± 0.0	1.7 ± 0.1
High CO <sub>2</sub> /low pH	1.9 ± 0.0	1.9 ± 0.0	1.9 ± 0.0	1.7 ± 0.1
High temp	1.8 ± 0.0	1.8 ± 0.0	1.8 ± 0.0	1.5 ± 0.1
High CO <sub>2</sub> /low pH + high temp	1.9 ± 0.0	1.9 ± 0.0	1.8 ± 0.0	1.5 ± 0.0

DIC (mmol L<sup>-1</sup>) values are mean ± SD.

**Table 2.** Dissolved inorganic phosphate ( $\text{PO}_4^{3-}$ ), nitrite + nitrate ( $\text{NO}_2^- + \text{NO}_3^-$ ) and ammonium ( $\text{NH}_4^+$ ) from each of the four pH and temperature conditions for units with and without *Dolichospermum* sp.

	$\text{PO}_4^{3-}$	$\text{NO}_2^- + \text{NO}_3^-$	$\text{NH}_4^+$
Without cyanobacteria			
Start	n.m.	n.m.	n.m.
Control	$9.2 \pm 0.0$	$51.3 \pm 2.1$	$0.9 \pm 0.2$
High $\text{CO}_2$ /low pH	$9.0 \pm 0.0$	$50.9 \pm 3.3$	$0.9 \pm 0.5$
High temp	$9.1 \pm 0.0$	$49.8 \pm 0.8$	$1.2 \pm 0.2$
High $\text{CO}_2$ /low pH + high temp	$9.0 \pm 0.1$	$51.0 \pm 1.1$	$1.2 \pm 0.1$
With cyanobacteria			
Start	$12.7 \pm 0.2$	$50.1 \pm 1.7$	$0.4 \pm 0.1$
Control	$10.5 \pm 0.5$	$52.4 \pm 2.0$	$0.3 \pm 0.2$
High $\text{CO}_2$ /low pH	$10.6 \pm 0.4$	$51.6 \pm 1.4$	$0.2 \pm 0.1$
High temp	$8.9 \pm 0.3$	$53.1 \pm 2.2$	$0.2 \pm 0.1$
High $\text{CO}_2$ /low pH + high temp	$8.9 \pm 0.3$	$52.2 \pm 1.6$	$0.2 \pm 0.2$

Start values are based on 4 replicates. All *Dolichospermum* sp. end samples are an average of 9 replicates  $\pm$  SD, whereas controls are an average of 3 replicates  $\pm$  SD. All concentrations are expressed as  $\mu\text{M}$ . n.m., not measured.

day 11 from all remaining nine replicate bottles of each treatment.

## Biomass and growth

Cyanobacterial biomass was assayed using POC, by filtering cells onto pre-combusted filters (Whatman GF/C, 4 h, 450°C, Little Chalfont, United Kingdom). The filters were dried for at least 24 h at 60°C and stored dry in a desiccator until analysis. POC was determined with a mass spectrometer (Elemental Combustion System CHNS-O 4010; Valencia, CA, United States), as described in Grasshoff (1976). Growth was calculated as  $(\ln \text{POC}_2 - \ln \text{POC}_1) / (t_2 - t_1)$  where  $\text{POC}_2$  and  $\text{POC}_1$  are the measured POC at time  $t_2$  and  $t_1$  (between day 0 and day 11).

## Toxin analysis

Intracellular microcystin levels were sampled by filtering the cultures onto a GF/F filter. All samples were frozen in  $-20^\circ\text{C}$  until further processing, according to instructions in Engström-Öst *et al.* (2011). Intracellular microcystin concentrations ( $\mu\text{g L}^{-1}$ ) were analyzed with enzyme-linked immunoabsorbant assay, using a microcystin plate kit (EnviroLogix; Portland, ME, USA), according to kit instructions. A negative control and a standard curve were measured accordingly. The toxin concentration was determined by dual reading from the absorbance at 450 nm and as reference 655 nm, measured with a photometer (Tecan infinite M200, Tecan Group Ltd, Männedorf, Switzerland), using Magellan software. Data are reported as microcystin-LR equivalents.

## Dissolved nutrients, inorganic carbon and pH

pH (pH-controller 7070/2, TUNZE) and temperature measurements were performed from each bottle at the start and at the end of the experiment (Table 1). After the dissolved inorganic carbon (DIC) measurement, pH was measured once more with pH meter Jenway (3510, calibrated with three buffers, Staffordshire, United Kingdom), when samples had

reached room temperature. DIC was measured according to Salonen (1981) from samples preserved in airtight bottles on ice ( $+3^\circ\text{C}$ ) and stored in darkness max 24 h until analysis (Table 3). Based on the measured DIC and the dissociation constant equations provided by Lueker *et al.* (2000), the concentrations of the three DIC forms ( $\text{CO}_2$ ,  $\text{HCO}_3^-$ ,  $\text{CO}_3^{2-}$ ) were calculated according to Zeebe and Wolf-Gladrow (2001) (Table 3). Samples for dissolved inorganic phosphate, ammonium and nitrite + nitrate were analyzed according to Grasshoff (1976) (Table 2).

## Oxidative status biomarkers

Oxidative status of the cyanobacteria was determined as intracellular soluble antioxidant capacity assayed by oxygen radical absorption capacity (ORAC) and lipid peroxidation, assayed as thiobarbituric acid reactive substances (TBARS). Samples were collected onto GF/C filters and were stored frozen in  $-80^\circ\text{C}$  until analysis. Each filter with cyanobacteria was homogenized in 1 mL of phosphate-buffered saline (PBS) for 4 min using FastPrep (Santa Ana, CA, USA) with cooling function and 100  $\mu\text{m}$  ceramic beads, centrifuged at 10 000  $g$  for 5 min at  $4^\circ\text{C}$  and supernatant was used to measure ORAC, TBARS and protein concentration. Then, supernatant were used for ORAC assay, 200  $\mu\text{L}$  for the lipid peroxidation assay and 50  $\mu\text{L}$  for protein determination.

Oxygen radical absorption capacity was measured in 200  $\mu\text{L}$  of the supernatant using oxidation of a fluorescent probe by peroxyl radicals by way of a hydrogen atom transfer process (Cao & Prior 1999; Prior *et al.* 2003) and a microplate reader (FLUORostar OPTIMA, BMG Labtech, Germany) set to 485 nm excitation/520 nm emission. The procedures were based on the modified ORAC<sub>FL</sub> method (Prior *et al.* 2003) using fluorescein (FL) as a fluorescent probe, 2,2'-Azobis (2-amidinopropane) dihydrochloride (AAPH) as a peroxyl radical generator, and Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid, a water soluble vitamin E) as a standard (6 to 50  $\mu\text{M}$  range). The final ORAC values were

**Table 3.** Contribution of different dissolved inorganic carbon forms; carbonate ( $\text{CO}_3^{2-}$ ), bicarbonate ( $\text{HCO}_3^-$ ), and carbon dioxide ( $\text{CO}_2$ ) given as  $\text{mmol L}^{-1}$ 

	$\text{CO}_3^{2-}$		$\text{HCO}_3^-$		$\text{CO}_2$	
	Start	End	Start	End	Start	End
Without cyanobacteria						
Control	0.04 ± 0.00	0.03 ± 0.00	1.74 ± 0.00	1.75 ± 0.02	0.03 ± 0.00	0.04 ± 0.00
High $\text{CO}_2$ /low pH	0.02 ± 0.00	0.02 ± 0.00	1.78 ± 0.02	1.77 ± 0.02	0.07 ± 0.00	0.07 ± 0.00
High temp	0.05 ± 0.01	0.04 ± 0.00	1.72 ± 0.01	1.73 ± 0.01	0.03 ± 0.00	0.03 ± 0.00
High $\text{CO}_2$ /low pH + high temp	0.02 ± 0.00	0.02 ± 0.00	1.80 ± 0.01	1.82 ± 0.01	0.07 ± 0.00	0.07 ± 0.00
With cyanobacteria						
Control	0.03 ± 0.00	0.24 ± 0.06	1.78 ± 0.02	1.41 ± 0.12	0.04 ± 0.00	0.00 ± 0.00
High $\text{CO}_2$ /low pH	0.02 ± 0.00	0.16 ± 0.06	1.79 ± 0.02	1.56 ± 0.11	0.09 ± 0.00	0.01 ± 0.01
High temp	0.05 ± 0.00	0.50 ± 0.06	1.69 ± 0.01	0.97 ± 0.09	0.03 ± 0.00	0.00 ± 0.00
High $\text{CO}_2$ /low pH + high temp	0.02 ± 0.00	0.47 ± 0.06	1.74 ± 0.02	1.05 ± 0.09	0.06 ± 0.00	0.00 ± 0.00

Values are shown units with and without *Dolichospermum* sp.

calculated using a linear regression between the concentration of Trolox standards and the net area under the FL decay curve for the standards.  $\text{ORAC}_{\text{FL}}$  values were expressed as  $\mu\text{M}$  trolox equivalent.

The lipid peroxidation assay was carried out using QuantiChrom TBARS Assay Kit (DTBA-100; BioAssay Systems, Hayward, CA, USA) as specified by the manufacturer with minor modifications. Standards were prepared by mixing 15  $\mu\text{L}$  of the 1.5 mM malondialdehyde (MDA) with 735  $\mu\text{L}$   $\text{dH}_2\text{O}$  and diluting to set a range 0 to 30  $\mu\text{M}$  MDA. To the 200  $\mu\text{L}$  of the supernatant, 400  $\mu\text{L}$  of trichloroacetic acid (TCA, 10%) were added, incubated on ice for 15 min and centrifuged at 14 000  $g$  for 15 min at 4°C. Two 200  $\mu\text{L}$  aliquots, one for sample and one for individual blank, were transferred to microcentrifuge tubes, and 200  $\mu\text{L}$  of the TBA Reagent were added to all tubes designated for samples and standards. Individual blanks were prepared by replacing the TBA Reagent with 3 mM HCl (pH 1.6). All tubes were then incubated at 100°C for 60 min. After cooling down to room temperature, samples and standards were loaded in duplicate to wells of a black flat bottom microplate (100  $\mu\text{L}$  well<sup>-1</sup>) and fluorescence was measured with the FLUOstar Optima microplate reader at excitation/emission wavelengths of 544/590 nm. The TBARS levels were expressed as nmol MDA per mg protein.

Water soluble protein concentration ( $\mu\text{g mL}^{-1}$ ) was measured with the bicinchoninic acid assay (Pierce, Rockford, IL, USA) with bovine serum albumin as standard following directions of the manufacturer for microtiter assays. In brief, 50  $\mu\text{L}$  of the supernatant were mixed with 250  $\mu\text{L}$  test kit reagents and incubated for 1 h at room temperature. The microplates were read at 550 nm using FLUOstar Optima microplate reader with absorbance configuration.

## Statistical analyses

A two-way analysis of variance (ANOVA) was used to test for differences between pH (normal and low) and temperature (normal and elevated) treatments and their interaction. Levene's test was used to assess homogeneity of variance. Spearman's rank correlation analysis was used to assess the relationship between toxin concentrations and growth rates


and partial correlation analysis to assess the relationship between ORAC and toxin concentrations. These statistical analyses were performed using SPSS 15.0 (SPSS, Chicago, IL, USA).

## RESULTS

The growth rates of the cyanobacterium varied between 0.06  $\text{g POC day}^{-1}$  in high  $\text{CO}_2$ /low pH and 0.11  $\text{g POC day}^{-1}$  in high  $\text{CO}_2$ /low pH + high temperature (Fig. 1a). Intracellular toxin concentrations varied between 3.6  $\text{mg g}^{-1}$  in high  $\text{CO}_2$ /low pH and 6.7  $\text{mg g}^{-1}$  in high temperature (Fig. 1b). ORAC was lowest in the control (105.2  $\mu\text{M}$  trolox equivalents), and highest in the high temp (132.2  $\mu\text{M}$  trolox equivalents) (Fig. 1c). Lipid peroxidation (measured as TBARS) varied between 0.64 mmol MDA  $\text{mg protein}^{-1}$  in the high temperature and 0.78 mmol MDA  $\text{mg protein}^{-1}$  in high  $\text{CO}_2$ /low pH + high temperature (Fig. 1d).

Growth rate was significantly positively affected by temperature, but not by  $\text{CO}_2$ /pH (Fig. 1a, Table 4). Intracellular microcystin concentrations per unit biomass were significantly positively affected by temperature (Fig. 1b, Table 4), but not by  $\text{CO}_2$ /pH. There was no interaction effect of  $\text{CO}_2$ /pH and temperature on microcystin concentrations (Fig. 1b, Table 4). Toxin concentrations were also significantly positively correlated with growth rates (Spearman's,  $R = 0.601$ ,  $n = 36$ ,  $P < 0.001$ ). There were some initial extracellular microcystin present,  $3.1 \pm 0.5 \mu\text{g L}^{-1}$ , but as the experiment progressed, the extracellular microcystin disappeared and was close to detection (data not shown).

Temperature,  $\text{CO}_2$ /pH and their interaction affected the oxidative status of the cells. ORAC responded positively to elevated temperature and  $\text{CO}_2$ /pH (Fig. 1c, Table 4). There was also an interaction effect of temperature and  $\text{CO}_2$ /pH on ORAC. In addition, there was a non-significant correlation between ORAC and toxin concentrations, with POC as a fixed variable (Partial correlation,  $R = 0.173$ ,  $n = 33$ ,  $P > 0.05$ ). TBARS was higher in high  $\text{CO}_2$ /low pH treatments (Fig. 1d). Moreover, there was a significant  $\text{CO}_2$ /pH  $\times$  temperature interaction (Table 4), suggesting increased lipid peroxidation in cells exposed to both increased  $\text{CO}_2$ /lowered pH and elevated temperature.



**Fig. 1.** Response of the cyanobacterium *Dolichospermum* sp. to high CO<sub>2</sub>/lowered pH (CO<sub>2</sub>/pH) and elevated temperature. (a) growth, (b) carbon normalized intracellular microcystin levels, (c) oxygen radical absorbance capacity (ORAC), and (d) lipid peroxidation, which was measured as thiobarbituric acid reactive substances (TBARS), at the end of the 11-day experiment. Mean ± SD, *n* = 9. Dashed lines represent average initial levels.

**Table 4.** Statistical results of two-way ANOVA, including *F*-values, *P*-values and degrees of freedom, assessing effects of temperature and CO<sub>2</sub>/pH

	<i>F</i> <sub>1,32</sub>	<i>P</i>
Growth		
Temp	74.768	<0.001
CO <sub>2</sub> /pH	0.25	0.876
Temp × CO <sub>2</sub> /pH	1.651	0.208
IC		
Temp	46.615	0.001
CO <sub>2</sub> /pH	7.308	0.132
Temp × CO <sub>2</sub> /pH	1.576	0.478
ORAC		
Temp	44.195	<0.001
CO <sub>2</sub> /pH	8.186	0.007
Temp × CO <sub>2</sub> /pH	29.526	<0.001
TBARS		
Temp	3.369	0.076
CO <sub>2</sub> /pH	43.380	<0.001
Temp × CO <sub>2</sub> /pH	6.530	0.016

IC, intracellular microcystin concentration; ORAC, oxygen radical absorbance capacity; TBARS, lipid peroxidation.

## DISCUSSION

Both growth rates and microcystin concentrations were significantly positively affected by temperature, regardless of CO<sub>2</sub>/pH. These results are supported by previous studies; Fu *et al.* (2007) reported higher cell division rates of picocyanobacteria *Synechococcus* in higher temperatures (+4°C). Similar results are shown also for other cyanobacteria: *Microcystis aeruginosa* Kützing (Imai *et al.* 2009; Bouchard & Purdie 2011; Dziallas & Grossart 2011), *Nodularia spumigena* (Lehtimäki *et al.* 1997) and *Anabaena* sp. currently *Dolichospermum*; Wagner & Adrian 2009; O'Neil *et al.* 2012). Increased growth due to warming is common among cyanobacteria (reviewed by O'Neil *et al.* 2012). Paerl and Scott (2010) suggest that higher temperatures will be a catalyst for future expanding cyanobacteria blooms in a warmer climate. Cyanobacteria generally exhibit optimal growth rates at relatively high temperatures and they also compete most effectively with eukaryotic primary producers at elevated temperatures (reviewed by Paerl & Huisman 2009).

Microcystin concentrations were significantly correlated with growth and this is supported in the literature (Orr & Jones 1998; Long *et al.* 2001). In the current study, microcystin concentrations and growth both increased significantly as a response to elevated temperature. Higher temperature may enhance cyanobacterial toxicity by stimulating toxin biosynthesis (El-Shehawy *et al.* 2011), but this process seems to be associated with growth. Dziallas and Grossart (2011) showed that the toxicity and also the toxic potential (i.e., the ratio of toxic vs. non-toxic cells) of *Microcystis aeruginosa* increased considerably in higher temperature. Further, a toxin-producing strain is more tolerant to oxidative stress than a non-toxic one (Dziallas & Grossart 2011), and it has been suggested that toxins protect the cell against stressors (Zilliges *et al.* 2011).

Effects of pH on cyanobacteria are also studied intensively and results commonly show increasing growth in systems with

continuous input of CO<sub>2</sub> (Hutchins *et al.* 2007; Levitan *et al.* 2010; Wannicke *et al.* 2012), with some exceptions (Czerny *et al.* 2009). In the current study, a single initial CO<sub>2</sub> addition and resulting change in pH did not change cyanobacterial growth significantly. Working with live, actively growing and CO<sub>2</sub> consuming phytoplankton in batch cultures, such as the *Dolichospermum* sp. culture in the current study, the initial CO<sub>2</sub> treatment is prone to changes, which was an expected outcome. We consider such a change realistic during a typical cyanobacteria bloom in the Baltic since the high production may quickly result in exhaustion of free CO<sub>2</sub> whereby pH may reach well beyond 9 (Hansen 2002; Brutemark *et al.* 2011). Thus, the difference between initial and end pH values reflect the growth of the cyanobacterium, similar to conditions prevailing during a cyanobacteria bloom in the Baltic Sea. Consequently, the high CO<sub>2</sub> demand of an algal bloom may result in carbon limitation. In our experiment, CO<sub>2</sub> depletion was evident in all treatments; and initial addition of CO<sub>2</sub> had no effect on growth, suggesting that *Dolichospermum* sp. was not carbon limited. Although not measured in the present study, it appears that *Dolichospermum* sp. is able to utilize HCO<sub>3</sub><sup>-</sup> and does not become carbon limited when CO<sub>2</sub> is diminished, a feature common among cyanobacteria (Kaplan *et al.* 1991; So & Espie 2005). In addition, dissolved nutrients (primarily nitrate and phosphate) were available in ample amounts throughout the experiment in all treatments, and the cyanobacterium was therefore not nutrient limited. CO<sub>2</sub>/pH did not significantly affect toxin concentrations in the current study, and only a few studies show the rise of toxin concentrations by increasing external CO<sub>2</sub> concentration (dinoflagellates, Fu *et al.* 2010; diatoms, Tatters *et al.* 2012).

Apart from influencing the water chemistry, pH can also have a direct effect on the cells by affecting metabolism and membrane transport, and thereby impairing cellular function or increasing cellular energy demand (Hinga 2002). Although CO<sub>2</sub>/pH in our study did not have an effect on growth, it did influence both antioxidant levels and lipid peroxidation. High temperature alone was beneficial as evidenced by high antioxidant levels. In contrast, when high temperature was combined with high CO<sub>2</sub>/low pH, the anti-oxidative level was most likely weakened by increased ROS production, which resulted in significant 20% lipid peroxidation increase (measured as TBARS). In cyanobacteria, respiration and photosynthesis are considered to be sources of ROS production (Latifi *et al.* 2009). When the balance between oxidant level and antioxidant production is lost, the organism faces oxidative stress that can generate damage on cell level (Latifi *et al.* 2009). Cell death due to oxidative stress is shown in *Microcystis aeruginosa* (Ross *et al.* 2006), and is suggested to be a major cause of bloom ending, shown in *Trichodesmium* (Berman-Frank *et al.* 2004). In our experiment *Dolichospermum* growth did not seem to be limited by carbon and we therefore suggest that pH, in combination with warming, have an oxidative effect on the cyanobacteria cells (Fig. 1). Both CO<sub>2</sub>/pH and temperature are stressors that can cause changes in the oxidative status of a photosynthetic organism (Bouchard & Purdie 2011). Bouchard and Purdie (2011) and references cited therein (Jochem 1999; Wu *et al.* 2008) suggest that cyanobacteria in general are tolerant of stress conditions as they are able to reduce their metabolic activity.

The role of microcystin has been debated, and recently it was suggested microcystin may act as defense against oxidative stress (Alexova *et al.* 2011; Dziallas & Grossart 2011; Phelan & Downing 2011). Consequently, we analyzed potential relationships between intracellular microcystin concentrations and ORAC, but as both were correlated to biomass, there was a risk of false relationship, and partial correlation analysis revealed that this was the case. Thus, the intracellular microcystin levels and ORAC are not directly associated, and there is, seemingly no relationship between cellular microcystin content and the oxidative status of the cells shown in the current study. However, to evaluate the response of *Dolichospermum* sp. to changes in CO<sub>2</sub>/pH and temperature we subjected the cyanobacterium to two CO<sub>2</sub>/pH levels (normal and high CO<sub>2</sub>/lowered pH) and two temperatures (normal and high) corresponding to CO<sub>2</sub>/pH changes that can occur in upwelling areas (Feely *et al.* 2008) as well as corresponding to temperature changes that are predicted for 2100 (Caldeira & Wickett 2003; IPCC 2007). The biological responses highlighted in our study may certainly differ in the presence of additional stressors and other cyanobacteria species/strains, but nonetheless, warming seems to benefit growth, toxicity, and antioxidant levels of *Dolichospermum* sp. When the cyanobacterium is exposed to the combination of elevated temperature and high CO<sub>2</sub>/lowered pH, it creates an imbalance between antioxidant levels and lipid peroxidation, indicating a shift of oxidative changes at cell level, while no simultaneous changes occurred in growth or toxicity. To conclude, it is suggested that escalating growth and toxicity of bloom-forming cyanobacteria as a response to warming have serious socio-economic implications (Paerl & Otten 2013), especially in heavily polluted basins such as the Baltic Sea.

## ACKNOWLEDGMENTS

The study was financially supported by the M and T. Nessling Foundation, the W and A. de Nottbeck Foundation, the R & D Institute Aronia, the Norden Havgruppe and the Academy of Finland (project numbers 125251, 255566, 276947). We thank Prof. K. Sivonen (University of Helsinki) for the BIR257 strain and U. Sjölund at Tvärminne Zoological Station for preparing the algal growth medium.

## REFERENCES

- Alexova, R., Haynes, P. A., Ferrari, B. C. and Neilan, B. A. 2011. Comparative protein expression in different strains of the bloom-forming cyanobacterium *Microcystis aeruginosa*. *Mol. Cell. Proteomics* **10**: M110.003749.
- Apel, K. and Hirt, H. 2004. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* **55**: 373–99.
- Barcelos e Ramos, J., Biswas, H., Schultz, K. G., LaRoche, J. and Riebesell, U. 2007. Effect of rising atmospheric carbon dioxide on the marine nitrogen fixer *Trichodesmium*. *Global Biogeochem. Cy.* **21**: GB2028.
- Barry, J. P., Widdicombe, S. and Hall-Spencer, J. M. 2011. Effects of ocean acidification on marine biodiversity and ecosystem function. In Gattuso, J.-P. and Hansson, L. (Eds) *Ocean Acidification*. Oxford University Press, Oxford, pp. 192–209.

- Berman-Frank, I., Biddle, K., Hamaraty, L. and Falkowski, P. G. 2004. The demise of the marine cyanobacterium, *Trichodesmium* spp., via an autocatalysed cell death pathway. *Limnol. Oceanogr.* **49**: 997–1005.
- Bouchard, J. N. and Purdie, D. A. 2011. Effect of elevated temperature, darkness, and hydrogen peroxide treatment on oxidative stress and cell death in the bloom-forming toxic cyanobacterium *Microcystis aeruginosa*. *J. Phycol.* **47**: 1316–25.
- Brutemark, A., Engström-Öst, J. and Vehmaa, A. 2011. Long-term monitoring data reveal pH dynamics, trends and variability in the western Gulf of Finland. *Oceanol. Hydrobiol. Stud.* **40**: 91–4.
- Caldeira, K. and Wickett, M. E. 2003. Anthropogenic carbon and ocean pH. *Nature* **425**: 365.
- Cao, G. and Prior, R. L. 1999. Measurement of oxygen radical absorbance capacity in biological samples. *Methods Enzymol.* **299**: 50–62.
- Collins, S. and Bell, G. 2004. Phenotypic consequences of 1,000 generations of selection at elevated CO<sub>2</sub> in a green alga. *Nature* **431**: 566–9.
- Czerny, J., Barcelos e Ramos, J. and Riebesell, U. 2009. Influence of elevated CO<sub>2</sub> concentrations on cell division and nitrogen fixation rates in the bloom-forming cyanobacterium *Nodularia spumigena*. *Biogeosciences* **6**: 1865–75.
- Dziallas, C. and Grossart, H. 2011. Increasing oxygen radicals and water temperature select for toxic *Microcystis* sp. *PLoS ONE* **6**: e25569.
- El-Shehawey, E., Gorokhova, E., Fernández-Piñas, F. and Campo, F. F. 2011. Global warming and hepatotoxin production by cyanobacteria: what can we learn from experiments? *Water Res.* **46**: 1420–9.
- Endres, S., Unger, J., Wannicke, N., Nausch, M., Voss, M. and Engel, A. 2013. Response of *Nodularia spumigena* to pCO<sub>2</sub> – part 2: exudation and extracellular enzyme activities. *Biogeosciences* **10**: 567–82.
- Engström-Öst, J., Repka, S. and Mikkonen, M. 2011. Interactions between plankton and cyanobacterium *Anabaena* with focus on salinity, growth and toxin production. *Harmful Algae* **10**: 530–5.
- Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D. and Hales, B. 2008. Evidence for upwelling of corrosive 'acidified' water on to the continental shelf. *Science* **320**: 1490–2.
- Feely, R. A., Alin, S. R., Newton, J. *et al.* 2010. The combined effects of ocean acidification, mixing, and respiration on pH and carbonate saturation in an urbanized estuary. *Estuar. Coast. Shelf Sci.* **88**: 442–9.
- Feng, Y., Warner, M. E., Zhang, Y. *et al.* 2008. Interactive effects of increased pCO<sub>2</sub>, temperature and irradiance on the marine coccolithophore *Emiliania huxleyi* (Prymnesiophyceae). *Eur. J. Phycol.* **43**: 87–98.
- Fewer, D. P., Köykkä, M., Halinen, K., Jokela, J., Lyra, C. and Sivonen, K. 2009. Culture-independent evidence for the persistent presence and genetic diversity of microcystin-producing *Anabaena* (Cyanobacteria) in the Gulf of Finland. *Environ. Microbiol.* **11**: 855–66.
- Fleming-Lehtinen, V., Laamanen, M., Kuosa, H., Haahti, H. and Olsonen, R. 2008. Long-term development of inorganic nutrients and chlorophyll *a* in the open northern Baltic Sea. *Ambio* **37**: 86–92.
- Fu, F.-X., Warner, M. E., Zhang, Y., Feng, Y. and Hutchins, D. A. 2007. Effects of increased temperature and CO<sub>2</sub> on photosynthesis, growth, and elemental ratios in marine *Synechococcus* and *Prochlorococcus* (cyanobacteria). *J. Phycol.* **43**: 485–96.
- Fu, F.-X., Zhang, Y., Warner, M. E., Feng, Y., Sun, J. and Hutchins, D. A. 2008. A comparison of future increased CO<sub>2</sub> and temperature effects on *Heterosigma akashiwo* and *Prorocentrum minimum*. *Harmful Algae* **7**: 76–90.
- Fu, F.-X., Place, A. R., Garcia, N. S. and Hutchins, D. A. 2010. CO<sub>2</sub> and phosphate availability control the toxicity of the harmful bloom dinoflagellates *Karlodinium veneficum*. *Aquat. Microb. Ecol.* **59**: 55–65.
- Grasshoff, K. 1976. *Methods of Seawater Analysis*. Verlag Chemie, Weinheim.
- Guillard, R. R. L. 1975. Culture of phytoplankton for feeding marine invertebrates. In Smith, W. L. and Chanley, M. H. (Eds) *Culture of Marine Invertebrate Animals*. Plenum Press, New York, pp. 29–60.
- Halinen, K., Jokela, J., Fewer, D. P., Wahlsten, M. and Sivonen, K. 2007. Direct evidence for production of microcystins by *Anabaena* strains from the Baltic Sea. *Appl. Environ. Microbiol.* **73**: 6543–50.
- Hansen, P. J. 2002. Effect of high pH on the growth and survival of marine phytoplankton; implications for species succession. *Aquat. Microb. Ecol.* **28**: 279–88.
- Hinga, K. R. 2002. Effects of pH on coastal marine phytoplankton. *Mar. Ecol. Prog. Ser.* **238**: 281–300.
- Hutchins, D. A., Fu, F.-X., Zhang, Y. *et al.* 2007. CO<sub>2</sub> control of *Trichodesmium* N<sub>2</sub> fixation, photosynthesis, growth rates, and elemental ratios: implications for past, present and future ocean biogeochemistry. *Limnol. Oceanogr.* **52**: 1293–304.
- Imai, H., Chang, K. H., Kusaba, M. and Nakano, S. 2009. Temperature-dependent dominance of *Microcystis* (Cyanophyceae) species: *M. aeruginosa* and *M. wesenbergii*. *J. Plankton Res.* **31**: 171–8.
- IPCC (Intergovernmental Panel on Climate Change). 2007. *Fourth assessment report*. [cited 28 August 2014]. Available from: <http://www.ipcc.ch>
- IPCC (Intergovernmental Panel on Climate Change). 2013. *Fifth assessment report*. [cited 28 August 2014]. Available from: <http://www.ipcc.ch>
- Jochem, F. J. 1999. Dark survival strategies in marine phytoplankton assessed by cytometric measurement of metabolic activity with fluorescein diacetate. *Mar. Biol.* **135**: 721–8.
- Kaplan, A., Schwarz, R., Lieman-Hurwitz, J. and Reinhold, L. 1991. Physiological and molecular aspects of the inorganic carbon-concentrating mechanism in Cyanobacteria. *Plant Physiol.* **97**: 851–5.
- Kótai, J. 1972. *Instructions for Preparation of Modified Nutrient Solution Z8 for Algae*. Publication B-11/69. Norwegian Institute for Water Research, Oslo.
- Kremp, A., Godhe, A., Egardt, J. *et al.* 2012. Intraspecific variability in the response of bloom-forming marine microalgae to changed climate conditions. *Ecol. Evol.* **2**: 1195–207.
- Latifi, A., Ruiz, M. and Zhang, C.-C. 2009. Oxidative stress in cyanobacteria. *FEMS Microbiol. Rev.* **33**: 258–78.
- Lehtimäki, J., Moisander, P., Sivonen, K. and Kononen, K. 1997. Growth, nitrogen fixation, and nodularin production by two Baltic Sea cyanobacteria. *Appl. Environ. Microbiol.* **63**: 1647–56.
- Levitan, O., Rosenberg, G., Setlik, I. *et al.* 2007. Elevated CO<sub>2</sub> enhances nitrogen fixation and growth in the marine cyanobacterium *Trichodesmium*. *Global Change Biol.* **13**: 531–8.
- Levitan, O., Sudhaus, S., LaRoche, J. and Berman-Frank, I. 2010. The influence of pCO<sub>2</sub> and temperature on gene expression of carbon and nitrogen pathways in *Trichodesmium* IMS101. *PLoS ONE* **5**: e15104.
- Long, B. M., Jones, G. J. and Orr, P. T. 2001. Cellular microcystin content in N-limited *Microcystis aeruginosa* can be predicted from growth rate. *Appl. Environ. Microbiol.* **67**: 278–83.
- Lueker, T. J., Dickson, A. G. and Keeling, C. D. 2000. Ocean pCO<sub>2</sub> calculated from dissolved inorganic carbon, alkalinity, and equations for K<sub>1</sub> and K<sub>2</sub>: validation based on laboratory measurements of CO<sub>2</sub> in gas and seawater at equilibrium. *Mar. Chem.* **70**: 105–19.
- Mazur-Marzec, H. and Pliński, M. 2009. Do toxic cyanobacteria blooms pose a threat to the Baltic ecosystem? *Oceanology* **51**: 293–319.

- Monaghan, P., Metcalfe, N. B. and Torres, R. 2009. Oxidative stress as a mediator of life history trade-offs: mechanisms, measurements and interpretation. *Ecol. Lett.* **12**: 75–92.
- O'Neil, J. M., Davis, T. W., Burford, M. A. and Gobler, C. J. 2012. The rise of harmful cyanobacteria blooms: the potential roles of eutrophication and climate change. *Harmful Algae* **14**: 313–34.
- Orr, P. T. and Jones, G. J. 1998. Relationship between microcystin production and cell division rates in nitrogen-limited *Microcystis aeruginosa* cultures. *Limnol. Oceanogr.* **43**: 1604–14.
- Paerl, H. W. and Huisman, J. 2009. Climate change: a catalyst for global expansion of harmful cyanobacterial blooms. *Environ. Microbiol. Rep.* **1**: 27–37.
- Paerl, H. W. and Otten, T. G. 2013. Harmful cyanobacterial blooms: causes, consequences, and controls. *Microb. Ecol.* **65**: 995–1010.
- Paerl, H. W. and Scott, J. T. 2010. Throwing fuel on the fire: synergistic effects of excessive nitrogen inputs and global warming on harmful algal blooms. *Environ. Sci. Technol.* **44**: 7756–8.
- Phelan, R. R. and Downing, T. G. 2011. A growth advantage for microcystin production by *Microcystis* PCC7806 under high light. *J. Phycol.* **47**: 1241–6.
- Prior, R. L., Hoang, H., Gu, L. *et al.* 2003. Assays for hydrophilic and lipophilic antioxidant capacity (oxygen radical absorbance capacity (ORAC<sub>FL</sub>)) of plasma and other biological and food samples. *J. Agric. Food Chem.* **51**: 3273–9.
- Riebesell, U., Fabry, V. J., Hansson, L. and Gattuso, J.-P. 2010. *Guide to Best Practices for Ocean Acidification Research and Data Reporting*. European Commission, Publications Office of the European Union, Luxembourg.
- Ross, C., Santiago-Vázquez, L. and Paul, V. 2006. Toxin release in response to oxidative stress and programmed cell death in the cyanobacterium *Microcystis aeruginosa*. *Aquat. Toxicol.* **78**: 66–73.
- Salonen, K. 1981. Rapid and precise determination of total inorganic carbon and some gases in aqueous solutions. *Water Res.* **15**: 403–6.
- Sellner, K. G. 1997. Physiology, ecology, and toxic properties of marine cyanobacteria blooms. *Limnol. Oceanogr.* **42**: 1089–104.
- Sivonen, K. and Jones, G. 1999. Cyanobacterial toxins. In Chorus, I. and Bartram, J. (Eds) *Toxic Cyanobacteria in Water – A Guide to Their Public Health Consequences, Monitoring and Management*. E & FN Spon, London, pp. 41–111.
- So, K.-C. and Espie, G. S. 2005. Cyanobacterial carbonic anhydrases. *Can. J. Bot.* **83**: 721–34.
- Sun, J., Hutchins, D. A., Feng, Y., Seubert, E. L., Caron, D. A. and Fu, F.-X. 2011. Effects of changing pCO<sub>2</sub> and phosphate availability on domoic acid production and physiology of the marine harmful bloom diatom *Pseudo-nitzschia multiseriata*. *Limnol. Oceanogr.* **56**: 829–40.
- Tatters, A. O., Fu, F.-X. and Hutchins, D. A. 2012. High CO<sub>2</sub> and silicate limitation synergistically increase the toxicity of *Pseudo-nitzschia fraudulenta*. *PLoS ONE* **7**: e32116.
- Unger, J., Endres, S., Wannicke, N. *et al.* 2013. Response of *Nodularia spumigena* to pCO<sub>2</sub> – part 3: turnover of phosphorus compounds. *Biogeosciences* **10**: 1483–99.
- Wacklin, P., Hoffmann, L. and Komárek, J. 2009. Nomenclatural validation of the genetically revised cyanobacterial genus *Dolichospermum* (Ralfs ex Bornet et Flahault) comb. nova. *Fottea* **9**: 59–64.
- Wagner, C. and Adrian, R. 2009. Cyanobacteria dominance: quantifying the effects of climate change. *Limnol. Oceanogr.* **54**: 2460–8.
- Wannicke, N., Endres, S., Engel, A. *et al.* 2012. Response of *Nodularia spumigena* to pCO<sub>2</sub> – part 1: growth, production and nitrogen cycling. *Biogeosciences* **9**: 2973–88.
- Wootton, J. T. and Pfister, C. A. 2012. Carbon system measurements and potential climatic drivers at a site of rapidly declining ocean pH. *PLoS ONE* **7**: e53396.
- Wu, Z., Song, L. and Li, R. 2008. Differential tolerances and responses to low temperature and darkness between waterbloom forming cyanobacterium *Microcystis* and a green alga *Scenedesmus*. *Hydrobiologia* **596**: 47–55.
- Zeebe, R. E. and Wolf-Gladrow, D. 2001. *CO<sub>2</sub> in Seawater: Equilibrium, Kinetics, Isotopes*. Elsevier Oceanography Series 65, Amsterdam.
- Zilliges, Y., Kehr, J.-C., Meissner, S. *et al.* 2011. The cyanobacterial hepatotoxin microcystin binds to proteins and increases the fitness of *Microcystis* under oxidative stress conditions. *PLoS ONE* **6**: e17615.