



This is an electronic reprint of the original article. This reprint may differ from the original in pagination and typographic detail.

Essential fish habitats (EFH): Conclusions from a workshop on the importance, mapping, monitoring, threats and conservation of coastal EFH in the Baltic Sea

Kraufvelin, Patrik; Pekcan-Hekim, Zeynep; Bergström, Ulf; Florin, Ann-Britt; Lehikoinen, Annukka; Mattila, Johanna; Olsson, Jens

Published: 01/01/2016

Link to publication

Please cite the original version:

Kraufvelin, P., Pekcan-Hekim, Z., Bergström, U., Florin, A.-B., Lehikoinen, A., Mattila, J., & Olsson, J. (2016). Essential fish habitats (EFH): Conclusions from a workshop on the importance, mapping, monitoring, threats and conservation of coastal EFH in the Baltic Sea. Nordic Council of Ministers. http://dx.doi.org/10.6027/TN2016-539

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.





Essential fish habitats (EFH)

Conclusions from a workshop on the importance, mapping, monitoring, threats and conservation of coastal EFH in the Baltic Sea





Essential fish habitats (EFH)

Conclusions from a workshop on the importance, mapping, monitoring, threats and conservation of coastal EFH in the Baltic Sea

Patrik Kraufvelin, Zeynep Pekcan-Hekim, Ulf Bergström, Ann-Britt Florin, Annukka Lehikoinen, Johanna Mattila and Jens Olsson

TemaNord 2016:539

Essential fish habitats (EFH)

Conclusions from a workshop on the importance, mapping, monitoring, threats and conservation of coastal EFH in the Baltic Sea Patrik Kraufvelin, Zeynep Pekcan-Hekim, Ulf Bergström, Ann-Britt Florin, Annukka Lehikoinen, Johanna Mattila and Jens Olsson

ISBN 978-92-893-4664-1 (PRINT) ISBN 978-92-893-4665-8 (PDF) ISBN 978-92-893-4666-5 (EPUB) http://dx.doi.org/10.6027/TN2016-539

TemaNord 2016:539 ISSN 0908-6692 © Nordic Council of Ministers 2016 Layout: Hanne Lebech Cover photo: Philippe Provençal

Print: Rosendahls-Schultz Grafisk Printed in Denmark



This publication has been published with financial support by the Nordic Council of Ministers. However, the contents of this publication do not necessarily reflect the views, policies or recommendations of the Nordic Council of Ministers.

www.norden.org/nordpub

Nordic co-operation

Nordic co-operation is one of the world's most extensive forms of regional collaboration, involving Denmark, Finland, Iceland, Norway, Sweden, and the Faroe Islands, Greenland, and Åland.

Nordic co-operation has firm traditions in politics, the economy, and culture. It plays an important role in European and international collaboration, and aims at creating a strong Nordic community in a strong Europe.

Nordic co-operation seeks to safeguard Nordic and regional interests and principles in the global community. Common Nordic values help the region solidify its position as one of the world's most innovative and competitive.

Nordic Council of Ministers Ved Stranden 18 DK-1061 Copenhagen K Phone (+45) 3396 0200

www.norden.org

Contents

Pre	eface	7	
Sui	Summary		
1.	General background1	3	
2.	Project objectives	7	
3.	Venue, participants and outline of the workshop19	9	
4.	Outcome of the workshop2	1	
	4.1 General evidence for relationships between habitat and	1	
	4.2 Manning and monitoring of EFH 2'	7	
	4.3 Conservation of and threats to EFH	3	
5.	General conclusions40	0	
6.	Future work44	2	
7.	Communication	4	
Ref	ferences4	6	
Sve	ensk sammanfattning54	4	
Ap	pendix 1 – Participants list58	8	
Ap	pendix 2 – Schedule of the workshop60	0	
Ap	 Appendix 3 – Abstracts from presentations		
	coastal habitats for fish stocks	3 1 2	
Ap	pendix 4 – Information from Baltic Sea countries about mapping and monitoring of essential fish habitats8	1	

Preface

Many coastal and offshore fish species in the Baltic Sea are strongly dependent on shallow and sheltered near-shore habitats for their spawning, nursery, feeding and migration. Still, the role of these essential fish habitats, EFH, for the development and support of fish stocks and communities has received relatively little attention. As EFH in the Baltic often are found in the same parts of the coastal zone that are highly utilized and valued by humans, they are subjected to many threats and therefore management needs are urgent. In addition, EFH provide and support important ecosystem services and are included in national and international agreements and legislative acts as the Baltic Sea Action Plan, Habitats Directive and Marine Strategy Framework Directive. Despite this, the conservation status of EFH is generally poor in the region.

As a result of these shortcomings and needs, the Nordic Council of Ministers financed a project including a workshop with Baltic Sea experts with the aim to review the importance of, protection of and threats to coastal EFH in the Baltic Sea. This report describes the results of the project, primarily focusing on the outcome of the workshop.

Figure 0: Group photo of the workshop participants



Summary

Many fish species in the Baltic Sea are highly dependent on shallow and sheltered coastal habitats that they use for spawning, nursery, feeding and migration. Still, the role of these essential fish habitats (EFH) for the development and support of fish stocks and communities has received relatively little attention, even though there is general consensus among scientists about their critical importance. Little is also known about the major threats to EFH, and their conservation status in different countries has previously not been reviewed. As EFH often are found in the same parts of the coastal zone that are also highly valued by humans, this gap in knowledge needs to be addressed. Hence, there is an urgent need to focus more thoroughly on the importance, mapping, monitoring and protection of EFH and also the driving factors and mechanisms behind the changes we observe in their status. Only this way, we will be able to predict and mitigate future effects of environmental change in these valuable habitats and to create adaptive management plans.

The main objectives of this project were to 1) organize a workshop for experts around the Baltic Sea on the importance, protection of and threats to coastal EFH (including an overview of the methods used for the mapping and monitoring of these habitats), and 2) based on the outcome of the workshop, produce a review paper in an international scientific journal about the state of the art of the subject for the Baltic Sea, including knowledge gaps and future research needs. Here, we report the results of the project, focusing on the outcome of the workshop.

From the workshop (organised during 2nd–4th June 2015 in Öregrund, Sweden) we conclude that there are only few quantitative studies available concerning the importance of EFH for fish stocks. This evidence is in turn quite complex and do not necessarily provide straightforward answers. Nevertheless, for some coastal species, indirect evidence exists and sufficient data are also available to carry out further quantitative analyses. More evidence on the role of EFH for fish production could potentially also be achieved using spatial and temporal data analyses, stage-structured modelling and otolith chemistry techniques. Based on qualitative results/analyses, it can be reasoned that EFH are very important and valuable for the provisioning of rich fish communities and for fish production. This conclusion is reached, despite the still quite low degree of targeted studies that are focusing explicitly on the role of the habitats and that are providing straight quantitative relationships. Most likely the importance of these habitats has been underestimated in the past and more studies could contribute to pinpoint their ecological importance.

For the monitoring and mapping aspects of EFH in Baltic Sea countries, a lot of data seems to be available. Different sampling methods are used for a wide range of both coastal and offshore species and life stages (from eggs, larvae, YOY (young of the year) to adult fish). The use of these data in producing habitat maps has for long been poor, but the situation is now improving rapidly in many countries as a result of national and regional underwater mapping and inventory programs. In this sense, not only habitat mapping and mapping of fish distribution (fish in different life stages) are of importance. Also the mapping of major threats, pressures and environmental background conditions should be performed. This would ensure maximum availability and optimized use of information necessary for efficient management and for the improvement of marine spatial planning.

The threats to and conservation status of EFH suggest urgent and diverse management solutions. Eutrophication, climate change, coastal construction and development, invasive species and fishery seem to constitute the major threats to the habitats. Among these threats, the physical pressures, including for example marine shipping/boat traffic and its associated infrastructure (like dredging), physical exploitation of shore areas and trawl fishery, tend to be more easily manageable. These activities disturb fish habitats both directly and indirectly, and are typically more serious to fish reproduction and juvenile stages. The conservation status of EFH is generally poor, mainly due to that fisheries management and nature conservation in the Baltic Sea region historically have been separated. Internationally, however, many marine ecological studies have shown how mutual benefits may be reached through an integrated management of fisheries and habitats. This gives EFH a central role in management, merging the interests of fisheries management and habitat protection, and simultaneously attracting a lot of scientific interest to associated research questions.

The results of the current project hence suggest that there are data for quantitative analyses to support the role of EFH for fish production, a potential to initiate, develop and synchronize future monitoring and mapping of the habitats, and that there is an increasing awareness for the protection and increased concern for the sustainability of these shallow coastal systems. The outcome of this project could serve as a basis for improving cooperation between Baltic Sea countries in this field, which in the long run could result in both harmonized monitoring and mapping methods of the EFH in the Baltic Sea and a strengthened management. The work also provides important input for developing indicators to assess the status of EFH and for the implementation of international agreements and legislative acts as the Baltic Sea Action Plan (BSAP), the Habitats Directive (HD) and the Marine Strategy Framework Directive (MSFD).

1. General background

The Baltic Sea is relatively shallow in relation to its size, and the coastal zone constitutes a large and ecologically important part of the system. The environmental status of many coastal areas of the Baltic Sea has declined during recent decades, partly as a result of increased eutrophication, but also due to climate change, coastal development and the introduction of non-indigenous species. Many studies report that evident changes in species composition of coastal fish and benthic communities have taken place during the past 30-40 years (e.g. Olsson et al. 2012, 2013a, Rousi et al. 2013, Snickars et al. 2015, Weigel et al. 2015). The habitat quality of shallow coastal ecosystems is also affected by non-indigenous species especially in inner coastal areas, since most of these species are originally spread by vessels (Katsenevakis et al. 2014, Ojaveer and Kotta 2014). In addition, evidence is accumulating for the occurrence of trophic cascades in the coastal system. In areas where populations of coastal predatory fish such as perch (Perca fluviatilis) and pike (Esox lucius) are weak, mesopredatory fish, such as the three-spined stickleback (Gasterosteus aculeatus), are present in high numbers (Eriksson et al. 2009, 2011, Bergström et al. 2015, Byström et al. 2015). These mesopredators can have substantial impacts on the community of invertebrate grazers, reducing their numbers and hence the grazing pressure on algae, ultimately leading to eutrophication symptoms and habitat changes with blooms of ephemeral filamentous algae (Korpinen and Jormalainen 2007, Baden et al. 2010, Sieben et al. 2011, Östman et al. 2016). Sticklebacks may also affect the egg and larval stages of many species of predatory fish negatively, further emphasizing this problem (Bergström et al. 2015; Byström et al. 2015).

An integrated management strategy which both includes fish and their preferred environments hence appears to be of key importance for combating eutrophication symptoms in coastal areas. Given the importance and vulnerability of coastal areas, a central focus on these parts of the Baltic Sea ecosystem in management would be pivotal for the future potential of the sea area to provide ecosystem goods and services (Ahtiainen and Öhman 2014, Sundblad and Bergström 2014, Sundblad *et al.* 2014, Bryhn *et al.* 2015). In order to halt the ongoing negative development and reverse the current unfavourable ecosystem status of the Baltic Sea, additional effective management actions are needed along with the current methods concentrating on reduction of nutrient loads. To support similar management measures, we need to build knowledge of the causal factors for ecosystem change and how these factors interact, and to monitor and assess the status of key ecosystem components such as for instance the availability and conditions of essential coastal habitats for fish, EFH.

EFH may be defined as environments necessary for any life-stage of a fish species and their importance can be assessed as the effects of changes in the quantity and/or quality of these habitats on populations or stocks in time or space (see e.g. Hansen and Snickars 2014, Sundblad *et al.* 2014). EFH thus includes nursery areas, feeding areas, spawning areas and migratory routes. The latter three habitats may be important for fisheries because of high catches or value per fishing effort. Therefore some habitats are important both for fish stocks and for fisheries, which creates challenges for their sustainable management. Despite the evident role of EFH for development of fish stocks, they have still in northern Europe received relatively little attention (but see Sundblad *et al.* 2014).

Many coastal fish species are highly dependent on shallow and sheltered coastal habitats, such as coastal wetlands, flads/lagoons/ bays/estuaries, seagrass beds, macroalgal beds, mussel beds, rocky bottoms (also artificial substrates), and unvegetated bottom habitats, for their reproduction (Airoldi and Beck 2007, Sundblad et al. 2011, Seitz et al. 2014). Coastal habitats are in addition also utilized as spawning and nursery areas of migratory marine species, such as herring and flatfishes. Sundblad et al. (2014) were among the first in the Baltic Sea to quantitatively demonstrate that the amount of suitable nursery habitat had a substantial impact on population densities of adult fish. In their study, the amount of essential coastal habitats explained about half of the variation in the adult population size. Despite the importance of EFH, their current status of protection is generally poor (Sundblad et al. 2011). Given that EFH are commonly found in parts of the coastal zone that are simultaneously subjected to many human activities, the exploitation rate of these spawning and nursery areas for coastal fish is high (Sundblad and Bergström 2014). Hence, there is an urgent need to focus on the role and protection of EFH, and on the driving factors and mechanisms behind the changes occurring in them. Apparently, there are also urgent needs for cross-sectoral management when it comes to EFH, i.e. fisheries management should consider environmental and habitat management and vice versa.

Despite that increased attention during recent years has been paid to characterizing, mapping and monitoring EFH around the Baltic Sea (HEL-COM 2012), adequate information is still lacking for most species in order to assess to which degree these coastal habitats limit population growth and production of fish. What we today also lack is a review of the work accomplished so far, including experiences of good practices and methods used. This gap in knowledge prevents quantitative comparisons of the results, and makes it difficult to draw general conclusions on the role of EFH in order to define the most urgent research needs to bring the work forward.

2. Project objectives

The main objectives of this project were to

- organize a workshop for experts around the Baltic Sea on the role and protection of EFH and for reviewing the methods used for mapping and monitoring these habitats
- based on the outcome of the workshop produce a review paper, about the state of the art of the subject in the Baltic Sea including knowledge gaps and future research needs, to be published in an international scientific journal
- initiate discussions and cooperation between Baltic Sea experts towards a common understanding for mapping, monitoring, protection and evaluation of EFH.

3. Venue, participants and outline of the workshop

The workshop was arranged in Öregrund in Sweden during 2nd–4th June 2015 and it gathered 30 participants from eight countries around the Baltic Sea: Finland, Estonia, Latvia, Lithuania, Poland, Germany, Denmark and Sweden. Of the Baltic Sea countries, participants were lacking only from Russia: (see Appendix 1. Participants list).

The workshop was structured around three central themes (Appendix 2. Agenda of the workshop):

- Theme 1 The role of essential coastal habitats for fish availability of quantitative evidence showing the importance of coastal habitats for fish stocks.
- Theme 2 How are essential coastal habitats for fish mapped and monitored? Methods, available maps, are there data for maps?
- Theme 3 Conservation of and threats to essential coastal habitats for fish.

4. Outcome of the workshop

In this section the primary outcomes of the workshop are presented and summarised per theme (see Appendix 3, Abstracts for presentations).

4.1 General evidence for relationships between habitat and population size

Theme 1 – The role of essential coastal habitats for fish – availability of quantitative evidence showing the importance of coastal habitats for fish stocks.

Day 1 was largely devoted to *Theme 1* and oral presentations were given by:

- Göran Sundblad, Sweden.
- Josianne Støttrup, Denmark.
- Henri Jokinen, Finland.
- Didzis Ustups, Latvia.
- Mehis Rohtla, Estonia.
- Timo Arula, Estonia.

In the following, the main background information and some case study examples as well as summaries and conclusions from the presentations within this theme are given.

A recent review by Seitz *et al.* (2014) has shown that 44% of all ICES species, i.e. the ones for which the International Council for Exploration of the Seas (ICES) do stock assessment (or give advice), utilize coastal habitats as spawning, feeding, nursery or migration areas and that these stocks contribute to 77% of the commercial landings of these species (Fig. 1). A limited habitat supply must therefore by some means control the size and dynamics of fish populations, but how can we find ways to quantify this? Species distribution modelling has emerged as a tool to map specific habitat requirements for different life stages of species with ontogenetic habitat shifts.

Basically, studies of fish populations appear to have evolved along two distinct paths, where one has centred on the dynamics of populations in

relation to their exploitation and another has focused on the static relationships between fish and their habitats. As quantitative evidence for habitat limitation is accumulating, from different areas and species through the use of different methods for quantification, there is now an unprecedented possibility, as well as a need, to integrate habitat in fisheries management and nature conservation. This is especially important in the light of efforts towards an ecosystem based approach to (fisheries) management (Appendix 3. Göran Sundblad's abstract).

Figure 1: Percentage (%) of ICES-advice fish species using coastal habitats for spawning, as nursery grounds, for feeding, and for migration



Source: Redrawn from data in Seitz et al. 2014.

4.1.1 Case studies on the importance of EFH

In Sweden and Finland, Sundblad *et al.* (2014) used species distribution modelling to relate distribution of nursery habitats for perch and pikeperch to the size of the adult populations in twelve archipelago areas. In this specific study, habitat availability explained almost half of the variation in population size, indicating a crucial role in limiting adult stock sizes. The relationships were, however, non-linear, suggesting that the effect of habitat loss or restoration would be largest in areas with little available habitat (Fig. 2). Especially important in this sense is that a map-based spatial approach yields quantitative links, which allows for map-based estimates of the regional distribution of large fish, scenario analyses, etc. (e.g. Bergström *et al.* 2013) (Appendix 3. Göran Sundblad's abstract).



Figure 2: Relationship between recruitment habitat in % and CPUE of a) perch and b) pikeperch

Source: Redrawn from data in Sundblad et al. 2014.

In Öresund, Denmark, multiple uses, such as commercial gillnet fishing, angling and sports diving as well as shipping, wind energy production and extraction of marine aggregates, may lead to conflicts between the fishing sector and between sectors impacting the sea floor and its habitats. Therefore, ecologically important habitats of selected fish species of commercial value were mapped during 2014–2015 based on descriptive information and fishermen interviews to visualize hot-spot fishery/fish habitat areas. Quantitative information from fishing surveys was, however, lacking and no attempt was made to relate the habitats to fish production (Appendix 3. Josianne Støttrup's abstract).

In Finland, the high diversity of habitats along the coast causes spatial overlap and mixing between different EFH, and makes it difficult to distinguish between specific effects of different EFH and to establish their relative roles for fish stocks. This may be exemplified by the flounder (*Platichthys flesus*) that is facing multiple pressures due to on-going large-scale ecosystem changes linked to e.g. eutrophication and climate change. Using available fishery-independent data on adult flounder as well as historical and present-state data on juvenile flatfish in nursery areas, a negative change over time may be demonstrated for both adults and juveniles, concurrent with increased coverage of filamentous algae in shallow areas (Jokinen *et al.* 2015, 2016). It is, however, difficult to find quantitative evidence on the role of the juvenile EFH for the adult flounder stock on the Finnish coast (Appendix 3. Henri Jokinen's abstract).

In Latvia, central Baltic Sea, pelagic and demersal spawning flounder (*P. flesus*) differ in their spawning habitat. In a recent study, it was examined whether the hydrological regime can explain fluctuations in early life

stages of pelagic spawning flounder (eggs, larvae, juveniles) over the past 30 years (Ustups *et al.* 2013). The hypothesis that the available reproductive volume (habitat), defined as the water column with dissolved oxygen larger than 1 ml/l and salinity between 10.6 and 12 PSU, affects the survival of flounder ichthyoplankton and determines recruitment success was evaluated. Both reproductive habitat volume and spawning stock biomass were significant factors determining flounder ichthyoplankton abundance. However, recruitment did not correlate with the supply of larvae, which could indicate that important bottlenecks are present in the shallow nearshore nursery grounds, utilized by both pelagic and demersal spawning flounder in the central Baltic Sea (Appendix 3. Didzis Ustups' abstract).

In Estonia, in the Väinameri Sea area, spawning habitat preference (i.e. fresh or brackish water) was investigated in brackish water populations of pike (Esox lucius), ide (Leuciscus idus) and burbot (Lota lota). Otolith Sr:Ca profiles were used to determine the hatching biome. Regarding adult pike, 90% hatched in freshwater and only 10% in brackish water (Rohtla et al. 2012), which may be compared with results from the Swedish waters, where 45% hatched in freshwater and 55% in brackish water (Engstedt et al. 2010). In another study from southern Sweden, no pike hatched in brackish water (Olsson et al. 2013b). These results suggest that brackish water spawning pike is becoming rarer, which could be due to deterioration of spawning grounds due to negative effects of eutrophication or overfishing. Regarding ide, 72% hatched in seasonal freshwater bays and only 28% in rivers (Rohtla et al. 2015). Regarding burbot, 96% hatched in fresh water and 4% showed signs of hatching in an environment with Sr:Ca slightly higher than the freshwater threshold, possibly in river mouths (Rohtla et al. 2014). In summary, also freshwater tributaries are of importance for coastal fish, but no analysis of the relationship between the rates of degradation of freshwater habitats and adult fish stocks has been undertaken (Appendix 3. Mehis Rohtla's abstract).

In another Estonian study from the NE part of Gulf of Riga (GoR), which has been the most important spawning and nursery ground for spring and autumn spawning herring populations and where larval herrings and the related environment has been studied regularly since 1947, larval herring individuals have been counted weekly. A new bottleneck may have been identified for spring spawning herring in form of temperatures exceeding the physiological thermal optimum for the survival of postflexion herring larvae in shallow estuarine habitats (Ojaveer *et al.* 2011, Arula *et al.* 2012a). Future climate warming indicates a risk for de-

creased survival of spring spawning larval herring, which may lead to reduced stocks of herring relying in recruitment in shallow coastal areas. With regard to Baltic autumn spawning herring, these landings constituted up to 47% in total herring landings in GoR, but have dropped drastically to <1% in the most recent decades (Arula *et al.* 2012b). In 2009– 2012, historically important spawning and nursery grounds for herring were visited and distribution and abundance of larvae were compared with historical data. It appeared that the same nursery grounds were in use and the number of larvae was comparable with the levels in the the1960s and 1970s before the herring stock collapsed. In conclusion, this nursery habitat does not seem to be limiting autumn spawning herring in the NE GoR (Appendix 3. Timo Arula's abstract).

4.1.2 Summary and conclusions of Theme 1

There are only few quantitative studies concerning the importance of EFH for fish stocks available in the Baltic Sea, and these do not provide straightforward answers. For most species, too little seems to be known in order to judge whether the coastal habitats are actually essential and limiting the production for fisheries (Seitz *et al.* 2014). Nevertheless, for some coastal species in some regions, indirect evidence exists, for example for flounder in the Gulf of Riga and Gulf of Finland. Curiously, there seems to be better evidence for non-migrating coastal species compared to migrating species. This could potentially be due to the conservative nature in habitat choice of non-migratory fish, or simply that it is easier to detect fish-habitat relationships in studies where many geographically restricted populations may be included. At some occasions, however, the necessary data for quantitative examination of the importance of EFH for fish stocks may already exist, but no analyses have yet been undertaken.

More evidence on the role of EFH could potentially be achieved using:

- Spatial approaches (e.g. assessing relationships between habitats of juveniles and adult fish to detect bottlenecks in early life stage).
- Temporal data analyses (e.g. assessing variability between years in success of different life stages).
- Stage-structured modelling (assessing habitat specific survival in stage-structured models) or
- Otolith chemistry techniques (comparing contribution of different habitats through "fingerprinting" the origin of fish).

On a general basis, it can be qualitatively reasoned that EFH are central for the provisioning of rich fish communities and for fish production, despite the so far low number of targeted studies focusing explicitly on the role of the habitats in the Baltic Sea. There may also be a great deal of indirect and qualitative evidence of the importance of EFH, and the role of freshwater habitats for coastal fish stocks should also be examined more closely. Most likely the importance of EFH has been underestimated in the past and more studies could contribute to pinpoint their ecological importance. A fruitful step forward could be to

- determine the current state of the art by forming country-wise roadmaps
- start-up investigations in countries where studies have yet not been conducted
- carefully looking into the methodologies and the needs and potentials for integration in a Baltic Sea-wide perspective.

Figure 3: The evening of Day 1 was closed with dinner and a boat trip in the Öregrund-Gräsö archipelago



4.2 Mapping and monitoring of EFH

Theme 2 – How are essential fish habitats mapped and monitored? Methods, available maps, are there data for maps?

There was only one scheduled presentation under *Theme 2* (See Appendix 2, Program) and that one was given by Meri Kallasvuo, Finland. Some material from presentations under *Theme 1* and *Theme 3* was, however, also utilised within this section of case studies.

4.2.1 Case studies

In Finland, systematic mapping of reproduction habitats for coastal fish has been conducted in the extensive national VELMU program during the years 2004-2015. VELMU covers the entire Finnish archipelago with 46,000 km of shoreline. Within VELMU, new field survey and mapping methods have been developed and tested and these approaches have resulted in modelled distribution maps of key reproduction habitats for the most important coastal fish species (Fig. 4, HELCOM 2012, see also Veneranta et al. 2011). These maps serve as basic references and allow visual and numerical comparisons of coastal areas. They also provide concrete support to fisheries and environmental management and they have been especially useful when planning coastal areas (MSP) and setting local or national fishing restrictions. Also, the developed cost-effective field sampling methods have been useful when more detailed surveys for local planning are conducted. Nevertheless, major challenges have arisen from the high annual variation in and insufficiency of field survey data. In addition, large archipelago areas and often strong environmental gradients require environmental data of high resolution and the poor availability of such data is a challenge. Efforts have anyway been put into building a VELMU online map service (www.http://paikkatieto.ymparisto.fi/velmu/map.htm) to openly share the data and knowledge in an easily accessible format (Appendix 3. Meri Kallasvuo's abstract).



Figure 4: A predictive map of roach reproduction areas in the western Gulf of Finland

Source: Finnish Game and Fisheries Research Institute (From HELCOM 2012).

In Estonia, coastal habitats have been mapped through exercises in which direct "point data" and indicator species serve as inputs. Mapping has been done for soft sediments, sandy bottoms, sandbanks, reefs, "tidal" mudflats and sandflats. In 2011–2014, spawning grounds for the Baltic herring around the Estonian coast have also been mapped using several methods and predictive distributional maps of herring spawning areas have been created (Appendix 3. Lauri Saks' abstract).

In Sweden, mapping of EFH has been initiated at various levels as an integrated part of the national management. Potential fish habitats may e.g. be mapped to indicate where one may expect certain life stages of specific fish to be distributed, i.e. where the environment is suitable for the species in question. Realized habitats, in turn, refer to habitats where the fish species actually are present, while effective habitats are those areas making a relatively high contribution to a fish population in comparison with adjacent areas. Species and habitat distribution maps may be used in marine spatial planning to protect sensitive areas from threats like shoreline development, dredging and boating (Fig. 5, Sundblad et al. 2011), and to evaluate and strengthen networks of marine protected areas (MPAs). To quantify threats to EFH, maps of the pressure variables, together with a mechanistic understanding of the effects of different threats on the habitats, are also needed. Having this information at hand, it is possible to perform scenario analyses to explore effects of different management regimes. Similar analyses may be useful in marine spatial

planning for efficient use of space (Appendix 3. Göran Sundblad's and Ulf Bergström's abstracts).



Figure 5: Map showing the representativity of fish recruitment habitats in Stockholm archipelago and in the Finnish Archipelago Sea

Source: From Sundblad et al. 2011.

In Germany, information about the eel stock has previously been collected to develop new management plans and these data have more recently been combined with sonar measurements to correlate the number of caught eel with the structure of the bottom (Ubl and Dorow 2015) (Appendix 3, Peter Möller's abstract).

In Lithuania, the bottom structure of coastal waters has been roughly mapped already several decades ago. More detailed maps have been produced during the recent decade as the result of various research projects. For instance, multibeam bathymetry maps and seabed profiles were developed to detect herring spawning beds and a recent LIFE-NATURE project used mapping of twaite shad and whitefish (*Coregonus lavaretus*) abundances in trawling during 1995–2010 to suggest EFH species-specifically (Appendix 3. Linas Ložys' abstract).

4.2.2 Questions regarding Theme 2

Within the *Theme 2*-session, each country was asked to provide answers to a list of questions regarding the topic. The following questions were asked:

- If there is monitoring of essential fish habitats in the coastal areas of your country, what habitats and what fish species are covered/in focus?
- What methods are used for the monitoring (give a brief description with reference to publications or methodological standards?
- What is the coverage of the monitoring in terms of time (are there time-series around) and what is the geographical coverage?
- What are the purposes/motivation/aims of the monitoring?
- Are there available maps (in which form) or available data on essential fish habitats? Please describe this rather detailed.
- Is the information from monitoring and/or mapping used in spatial planning?
- In your country or in general, what do you feel is the future for monitoring and mapping of EFH (especially in light of the upcoming EU directive on marine spatial planning)? What indications are there from managers, stakeholders and politicians? Is there an increasing, stable or decreasing demand for this kind of information?

The results from this exercise can be found in Appendix 4 and they are also briefly summarised below.

4.2.3 Summary and conclusions of Theme 2

In summarising the presentations under *Theme 2* and the information collected within Appendix 4, there are considerable differences in mapping and monitoring of EFH between Baltic Sea countries with regard to fish species targeted, habitat characterization, methods used, spatial and temporal coverage, purpose/motivation and aims of the monitoring, as well as the availability of maps and data.

Regarding targeted fish species, some countries (e.g. Latvia, Lithuania) have mainly focused on specific species within targeted sampling programs, while others have sampled all or most occurring fish species (e.g. Estonia, Poland). Regarding habitats, some countries have very detailed information on sampling and characterisation of the habitats (e.g. Estonia, Finland, Poland, Sweden), while there is yet quite little information from other countries or the information has been gathered within other specific sampling programs (e.g. Denmark, Germany, Latvia, Lithuania).

The methods for fish data collection include standardised monitoring routines for coastal areas of the northern Baltic Sea, fishers interviews (Denmark and Sweden mainly), multimesh gillnets (all countries), beach seines (all countries), white plates (Finland), detonations (Finland, Sweden), dipnets, pushnets, SCUBA-diving, underwater video cameras (Germany, Latvia), hydroacoustic surveys or sonars (Germany, Latvia, Lithuania, Sweden), eel sampling system (Germany, Latvia, Sweden), trawl nets (Germany, Lithuania), tags (inside and outside), electro fishing (Germany, Poland), egg sampling (Germany, Latvia, Sweden), fish traps, multibeam bathymetry (Latvia and Lithuania) and side-scan sonar, benthic roe samples, bongo net hauls, neuston net hauls and fyke nets (Estonia, Finland, Poland, Sweden).

Habitat information has been obtained by extensive field surveys including remote sensing, SCUBA-diving, drop-videos and grab samples, and habitat maps have been produced in combination with statistical modelling, or, as a first and simpler approach, by interview studies.

Regarding fish species, there is information on coastal and migratory marine species, anadromous and catadromous species, as well as freshwater species with a coastal distribution in the Baltic Sea. The life stages targeted range from eggs, larvae, YOY (young of the year) to adult fish.

Regarding mapping methods, the following techniques have been used:

- Overlay techniques and simple visualisation techniques using geographical information systems (GIS) to plot spatial distributions.
- Interpolation techniques using GIS.
- Statistical modelling using point data on fish and environmental variables and subsequent map predictions using full-coverage maps of predictor variables and GIS. Statistical methods used are different regression techniques (e.g. GLM, GAM), classification techniques (e.g. Random forest), Gaussian process modelling and machine learning methods (e.g. maximum entropy modelling).
- Comprehensive interview studies, where spawning areas pointed out by fishers are delineated using GIS.

When it comes to spatial coverage of the sampling, most areas of the Baltic Sea seem to be covered in Appendix 4, except for the Russian areas (both the inner Gulf of Finland and Kaliningrad area), Western Baltic Sea coast of Germany and part of the Polish coast. Regarding temporal coverage, most information is from year 2000 and onwards, but occasionally there is also information already from the 1990s (in Poland even from the 1970s and 1980s).

The purposes of sampling span from coastal fish monitoring, inventories, mapping, stock assessments, research, monitoring of invasive species to marine spatial planning. With regard to marine spatial planning, mainly Finland and Latvia claimed to carry out sampling specifically for this purpose.



Figure 6: View of one of the workshop sessions within Theme 3 during Day 2

The use of habitat maps for management or for marine spatial planning is not very frequent or efficient yet, but the situation seems to be improving rapidly in many countries thanks to national and regional underwater mapping and inventory projects and programs. In Finland, for instance, there are now comprehensive maps on e.g. prediction maps for fish, benthic algae and macrofauna available from the national VELMU program (www.http://paikkatieto.ymparisto.fi/velmu/map.htm). In Germany, the monitored and mapped information is used to declare particular areas protected or even closed, like spawning grounds and also for planning marine construction projects, in particular the choice of areas suitable for wind mill parks (controversial discussions are currently going on). According to the participating experts in the workshop, there is clearly an increasing demand for monitoring and mapping of EFH and quite a lot of activities are going on. Despite the recent progress, however, there are still needs to improve mapping and characterization of current distribution of key habitats and fish production areas, and their internal relationships also need to be better quantified in many areas (Sundblad *et al.* 2011). In this sense, not only habitat mapping and mapping of occurrence and abundance of different species and life stages of fish species are of importance. Also mapping of major pressures and environmental background conditions are needed in order to make maximum use of all information that may be needed for efficient management and improved marine spatial planning.

The most important notation regarding this theme is, however, that some useful information on EFH seems to be available from all eight participating Baltic Sea countries, despite the scatter in applied methodologies and the level of data resolution. One way forward to improve harmonization of the methods and data between countries is to use the gathered information, describe the various approaches used and their applicability and further try to evaluate which ones are the best for specific conditions and situations as summarized in Appendix 4.

4.3 Conservation of and threats to EFH

Theme 3 – Conservation of and threats to essential coastal habitats for fish. Day 2 was largely devoted to *Theme 3* and oral presentations were given by:

- Patrik Kraufvelin, Finland/Sweden.
- Ulf Bergström, Sweden.
- Antti Lappalainen, Finland.
- Elliot John Brown, Denmark.
- Linas Ložys, Lithuania.
- Martin Snickars, Finland.
- Peter Möller, Germany.
- Zusana Celmer, Poland.

(Abstracts from the presentations can be found in Appendix 3).

In addition to presentations, each country was asked to provide answers to the following list of questions:
- What are the major threats to essential fish habitats (EFH) in coastal areas in your country?
- Is fisheries management (FM) and nature conservation (NC) split between authorities in your country? Is the situation changing?
- Is fish habitat protection an objective in NC?
- Is fish habitat protection an objective in FM?
- Are authorities aware of the importance of EFH? What about the public? Is the situation changing?
- Are fish habitat maps used in MSP (nationally or regionally)?
- Is maintenance/restoration of fish stocks an objective in NC?

4.3.1 Threats to EFH

Eutrophication, climate change, coastal construction and development, invasive species and fisheries have in general been acknowledged as the major threats to EFH (Turner et al. 1999, Jackson et al. 2001, Lotze et al. 2006, Orth et al. 2006). Among the abovementioned threats, physical pressures, including marine shipping and boat traffic with the infrastructure required (including dredging), coastal engineering measures in bays or on shores and trawl fishery (Kraufvelin et al. in prep.), are the ones most easily managed. Most of the stressors are more serious for fish reproduction and especially for juvenile stages, as these life stages are more strongly dependent on specific habitats and at the same time less mobile than the adult fish. It is noteworthy that the total impact of small boat traffic may be worse than the effect of ship traffic due to the high frequency of the former in shallow sheltered areas with fine sediment bottoms that are not naturally subjected to wave action, and because small boat traffic is most intense during the biologically most active and sensitive summer period (Sandström et al. 2005). To quantify threats to EFH, maps of the pressure variables, together with a mechanistic understanding of the effects of different threats on different habitats, are needed in addition to habitat maps (Appendix 3. Patrik Kraufvelin's abstract).

There were differences in the opinions of the experts regarding the major threats on the EHF between the Baltic Sea countries (Table 1). Likely explanations for the differences identified include for example geographic location, human development and economic status, type of coastline (archipelago versus open) and historical land use. The list only reflects the views of the participants in the workshop and not the official national views, which needs to be kept in mind. The following major threats were listed (number of countries that mentioned the threat/total number of countries):

- Eutrophication (7/8).
- Climate change (5/8).
- Coastal construction and development, including dredging (5/8).
- Invasive species (5/8).
- Fishing/fishery (legal/illegal) (3/8).

Marine litter, erosion, pollution, tourism, mussel fishery, marine extractions and wetland drainage are threats that were only mentioned by 1–2 countries each (Table 1).

Table 1: Summary of perceived threats to EFH

Country	Most important threat(s)	Important threat(s)	Other threats
Sweden	Eutrophication	Coastal construction and de- velopment	Fishing, climate change, inva- sive species, marine litter
Finland	Eutrophication and climate change	Coastal construction	Dredging
Estonia	Eutrophication (also effects from the past)	Dams (mainly past threats)	Climate change, draining of wetlands, marine extraction (sand removal)
Latvia	Erosion (wave impacts)	Invasive species	Eutrophication
Lithuania	Invasive species	Fishery	Cormorants, climate change
Poland	Construction and shipping	Tourism	Illegal fisheries, eutrophica- tion, pollution, invasive species
Germany	Eutrophication (internal load- ing from sediments)	Coastal construction (mari- nas) and gravel dredging	Tourism
Denmark	Eutrophication and climate change	Coastal construction	Cormorants (inner waters) and seals, invasive species, mussel fishery, dredging, ma- rine extraction

4.3.2 Current protection of EFH

The protection of EFH in the Baltic Sea varies greatly between countries and between types of habitats. In Finland for instance, shallow sandy beaches are protected by the Nature Conservation Act, small pristine flads (<10 ha) are protected by the Water Act and flads in Natura 2000 areas are protected by the Habitat Directive. There are also private conservation areas and fishing restrictions during spawning time. Inner bays and estuaries are generally not very strictly protected. In Estonia, large parts of the coastal area are relatively well protected, mostly as Natura 2000 areas, which have different restrictions that apply under various circumstances. For example, fishing restrictions for different areas can now be easily found at: http://pump.regio.ee/kalandus/public/ (Appendix 3. Antti Lappalainen's and Lauri Saks' abstracts).

In order to attain a better protection of EFH, the value of habitats in supporting fisheries must, however, be investigated in a broader context emphasizing also the values of the habitats in providing ecosystem services. These values may thus include producing fish for commercial and recreational fisheries, aquaculture and biological regulation (i.e. regulation of eutrophication symptoms through top-down control of filamentous algae). The habitats are also of importance for protection against erosion, as nutrient filters, and for human recreation. Natural scientists should therefore consider all the ways in which coastal fish habitats provide value to society and use these as examples when communicating the need for the protection of EFH and their sustainability. In this context, the need for protection of EFH from diverse pressures and what level of sustainable use of EFH that can be permitted should also be clearly stated (Appendix 3. Elliot John Brown's abstract).

4.3.3 Current management of EFH

Management of fisheries and nature conservation have historically been separated in the Baltic Sea region, and the awareness of potential synergistic effects between the two has been low. Recently, several marine ecological studies have shown how mutual benefits may be reached through an integrated management of fisheries and the environment (Pikitch *et al.* 2004, Möllmann *et al.* 2014, Seitz *et al.* 2014, Sundblad *et al.* 2014). In contrast to the situation in northern Europe, there is much stronger focus on EFH in the research and management in e.g. the USA and Canada. In these countries, management of EFH is regulated by legal acts.

Healthy vegetated habitats are needed for the reproduction of many coastal fishes (Rangeley and Kramer 1995, Stål *et al.* 2007, Snickars *et al.* 2010, Seitz *et al.* 2014), while strong populations of predatory fish may support habitat-forming vegetation through maintaining trophic cascades (Norderhaug *et al.* 2005, Moksnes *et al.* 2008, Newcombe and Taylor 2010, Eriksson *et al.* 2011, Svensson *et al.* 2012). To increase the awareness of the benefits of integrating management of fisheries and habitat, the scientific community could contribute in many ways. This could, for example, be done by demonstrating ecological synergies that may be

achieved by protecting EFH, developing methods for and carrying out large-scale mapping of EFH, quantifying the effects of different threats to EFH (see e.g. Fig. 7), and by communicating the importance of the habitats. The latter might be mediated through valuation of ecosystem services, such as food, recreation and biological regulation (Ahtiainen and Öhman 2014, Bryhn *et al.* 2015). An efficient way to increase awareness of and willingness to protect these sensitive coastal environments may also be through revealing their visual beauty with photos and videos (Appendix 3. Ulf Bergström's abstract).



Figure 7: Predicted effects on the distribution of recruitment areas (EFH) of perch and pikeperch as a response to changes in water transparency according to a set of eutrophication scenarios

Source: From Bergström et al. 2013.

In the Baltic Sea area, fisheries management (FM) and nature conservation (NC) are split among authorities in most countries with Sweden and partly Germany as exceptions. In Sweden, FM and NC are placed within the same authorities both regionally (county boards) and nationally (since 2011: Swedish Agency for Water Management), although they be-

Note: Curves show percentage change in areal cover with changes in Secchi depth, where numbers on x-axis denote % deviation from current Secchi depth level. Dotted lines show standard errors of predictions from three separate modelling methods. The arrows indicate, from left to right, the Secchi depth changes according to the scenarios business-as-usual (Scen. 1), Baltic Sea Action Plan (Scen. 2) and reference levels (Scen. 3).

long to different ministries. In Germany, FM and NC are split on the federal level with the Federal Ministry of Food and Agriculture taking care of fisheries management and the Federal Ministry for the Environment, Nature Conservation, Building and Nuclear Safety taking care of nature conservation. In Mecklenburg-Western Pomerania, however, FM and NC are both under the same roof, Ministry of Agriculture, Environment and Consumer Protection. In Denmark and Finland, where FM and NC are split between authorities, there are plans for fusion of ministries in the near future. In the other four countries, where FM and NC are split, i.e. in Estonia, Latvia, Lithuania and Poland, no changes are being scheduled.

Regarding the question if fish habitat protection is an objective of nature conservation or an objective of fisheries management, there are slight differences among the countries. Generally, fish habitat protection seems to be an objective within both FM and NC and this is the case for Denmark, Germany, Lithuania and Sweden (but for Sweden to a lesser extent in FM). In Poland, fish habitat protection clearly belongs to FM, while in Estonia and Latvia, fish habitat protection is mainly a task for NC. In Finland, there has not been very much focus on fish habitat protection in neither NC nor in FM, although the situation seems to be improving.

Generally, there is a need for more awareness of the importance of EFH among managers, politicians and the public, although the awareness seems to be increasing especially within authorities. Regarding the use of fish habitat maps in marine spatial planning, the information is used for these purposes in some countries and for some cases, but still, more advice on how to use the information and how to include it in marine spatial planning is needed. Finally, when it comes to maintenance and restoration issues of fish stocks as an objective in nature conservation, there is specific focus on salmonids, species included in the Habitats Directive, and also species that are red-listed. According to many experts, coastal habitats (benthic habitats – benthic fauna, macrophytes) in many countries have also been a focus of national conservation issues. In cases where there is focus on fish, it is, however, mainly for commercial and threatened species.

In summarising these results, it appears that the management of and the level of information and knowledge on EFH varies among countries. Awareness of the importance of EFH does, however, seem to be increasing among authorities. In contrast, the awareness of the importance of EFH among the general public, and more specifically among fishermen, is rather good, but knowledge on the threats to these habitats is considerably low. In some countries, the public does not seem to know much about these topics, which has also previously been concluded in other studies (Lotze 2004). In this sense, a great deal of outreach work is needed to increase awareness of the importance of EFH.

5. General conclusions

Available information suggests that fish habitats are essential for fish production and for the provisioning of rich fish communities in the Baltic Sea. Most likely the importance of EFH has been underestimated in the past and more studies could contribute to pinpoint their ecological importance. Sometimes, indirect evidence exists or the necessary data for quantitative examination of the importance of EFH for fish stocks may already be available, although no analyses have yet been undertaken.

Mapping and monitoring activities are generally well developed in different Baltic Sea countries with regards to a multitude of fish species including also different life stages. The methods used are diverse and differ between countries, species and life stages, partly by necessity but also partly by tradition. One way to initiate a harmonization of data collection methods and datasets would be to use the gathered information within this project and describe the various approaches used and their applicability and further try to evaluate which ones seem to be the best for specific conditions and situations. The use of habitat maps for EFH is generally rather poor, but seems to be rapidly improving in many countries as a result of national and regional underwater mapping and inventory programs. However, not only habitat mapping and mapping of fish occurrence or abundance are of importance, but also mapping of major pressures and environmental background conditions are needed in order to make maximum use of all information needed for efficient management and improved marine spatial planning.

With regard to the threats to EFH, eutrophication, climate change, coastal construction and development, invasive species and fishery were perceived as the most serious ones by the experts that participated in the workshop. Out of these, several are manageable in turn suggesting that the situation could be improved.

Management of fisheries and nature conservation have historically been separated in the Baltic Sea region, and the awareness of potential synergistic effects between the two areas has been low. However, the situation appears to be improving, and increased communication on the importance of the habitats is one way to mediate further progress. There is a great need for more knowledge about EFH and also for improved protection of the habitats. Currently the status of protection is rather low, primarily due to lack of research and information about the importance of EFH, as well as due to lack of maps depicting the spatial distribution of EFH to be used in spatial planning and other management purposes. In this respect, a great deal of outreach work should specifically be targeted towards managers, authorities, politicians, and users of the sea, such as fishers – but also towards media, teachers and the general public.

6. Future work

This project has brought about many suggestions for future work and cooperation. An important first step is that cooperation and discussions have now been initiated with regard to EFH all around the Baltic Sea. The long-term results of the work done so far will hopefully be the initiation of a Baltic-wide network of experts working on mapping and monitoring of EFH. Hopefully, this will create a common understanding of the role of EFH, how to investigate their importance for fish production, how to map and monitor them, and finally how to manage the threats to them and their status. Further studies seem to be especially urgent within the field of attaining quantitative data for the value of EFH for fish production including defining the key habitats for protection and restoration efforts. To improve the integration of habitat quality in fish stock assessment and ecosystem-based fishery management is also warranted (Seitz et al. 2014, Sundblad et al. 2014). An important part of this work should be formed by additional analyses on existing data as a lot of the needed information already seems to be available in many countries. Thus, the usage of some type of a meta-analytical approach could be worth considering. Common projects on e.g. mapping and interview studies could furthermore be other fruitful ways forward. There are also evident needs for projects to communicate the information (VELMU example), e.g. outreach efforts with needs to produce and assemble photos, maps, graphs, and education material for schools, teachers, managers and policy makers. In order to succeed with all these purposes, devoted projects focusing on EFH will be needed.

7. Communication

Besides this report, the results from this project will also be communicated as a review paper that will be submitted to an international scientific journal. In addition, the outcome of the project will be presented to national and local managers and stakeholders working with coastal ecosystem and fisheries issues in the countries participating in the workshop. For the international perspective, the results of the project was presented to relevant HELCOM groups HELCOM FISH PRO II (HELCOM expert group on coastal fish; presentation held at the HELCOM FISH PRO II meeting in Riga 16–18th of February 2016) and HELCOM FISH group (presentation held at the HELCOM FISH meeting in Gothenburg 11–12th of May 2016). The latter group focuses on bridging knowledge between nature conservation and fisheries management. The work will also provide important input for developing indicators and implementation of international agreements and legislative acts as the Baltic Sea Action Plan (BSAP, HEL-COM 2007), the Habitat Directive (HD) and the Marine Strategy Framework Directive (MSFD, Anon 2008). The results will further be applied for guidance on current and future methods for how to map and monitor EFH in the Baltic Sea.

References

- Ahtiainen, H. & Öhman, M.C. (2014). *Ecosystem services in the Baltic Sea: valuation of marine and coastal ecosystem services in the Baltic Sea*. TemaNord 2014:563. Nordic Council of Ministers, Copenhagen, 74 pp. http://dx.doi.org/10.6027/TN2014-563
- Airoldi, L. & Beck, M.W. (2007). Loss, status and trends for coastal marine habitats of Europe. *Oceanography Marine Biology: an Annual Review* 45, 345–405.
- Anon (2008). Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy. *Official Journal of the European Union* L164, 19–40.
- Arula, T., Kotta, J., Lankov, A., Simm, M. & Põlme, S. (2012a). Diet composition and feeding activity of larval spring-spawning herring: Importance of environmental variability. *Journal of Sea Research* 68, 33–40. http://dx.doi.org/10.1016/j.seares.2011.12.003
- Arula, T., Ojaveer, H. & Shpilev, H. (2012b). Individual fecundity of the autumn spawning Baltic herring *Clupea harengus membras* L. *Estonian Journal of Ecology* 61, 119–134. http://dx.doi.org/10.3176/eco.2012.2.04
- Baden, S., Boström, C., Tobiasson, S., Arponen, H., & Moksnes, P.O. (2010). Relative importance of trophic interactions and nutrient enrichment in seagrass ecosystems: A broad-scale field experiment in the Baltic-Skagerrak area. *Limnology and Oceanography* 55, 1435–1448. http://dx.doi.org/10.4319/lo.2010.55.3.1435
- Bergström, U., Sundblad, G., Downie, A.-L., Snickars, M., Boström, C., & Lindegarth, M. (2013). Evaluating eutrophication management scenarios in the Baltic Sea using species distribution modelling. *Journal of Applied Ecology* 50, 680–690. http://dx.doi.org/10.1111/1365-2664.12083
- Bergström, U., Olsson, J., Casini, M., Eriksson, B. K., Fredriksson, R., Wennhage, H., & Appelberg, M. (2015). Stickleback increase in the Baltic Sea A thorny issue for coastal predatory fish. *Estuarine Coastal and Shelf Science* 163, 134–142. http://dx.doi.org/10.1016/j.ecss.2015.06.017
- BONUS INSPIRE manual. http://www.bonus-inspire.org/
- Bryhn, A., Lindegarth, M., Bergström, L., & Bergström, U. (2015). Ekosystemtjänster från svenska hav. Status och påverkansfaktorer. Havs- och vattenmyndighetens rapport 2015:12. (In Swedish).
- Byström, P., Bergström, U., Hjälten, A., Ståhl, S., Jonsson, D., & Olsson, J. (2015). Declining coastal piscivore populations in the Baltic Sea: Where and when do sticklebacks matter? *Ambio* 44, 462–471. http://dx.doi.org/10.1007/s13280-015-0665-5
- Engstedt, O., Stenroth, P., Larsson, P., Ljunggren, L., & Elfman, M. (2010). Assessment of natal origin of pike (*Esox lucius*) in the Baltic Sea using Sr:Ca in otoliths. *Environmental Biology of Fishes* 89, 547–555. http://dx.doi.org/10.1007/s10641-010-9686-x
- Eriksson, B.K., Ljunggren, L., Sandström, A., Johansson, G., Mattila, J., Rubach, A., Råberg, S., & Snickars, M. (2009). Declines in predatory fish promote bloom-forming macroalgae. *Ecological Applications* 19, 1975–1988. http://dx.doi.org/10.1890/ 08-0964.1

- Eriksson B.K., Sieben, K., Eklöf, J., Ljunggren, L., Olsson, J., Casini, M., & Bergström, U. (2011). Effects of altered offshore food webs on coastal ecosystems emphasize the need for cross-ecosystem management. *Ambio* 40, 786–797. http://dx.doi.org/10.1007/s13280-011-0158-0
- Florin, A.-B., Sundblad, G., & Bergström, U. 2009. Characterisation of juvenile flatfish habitats in the Baltic Sea. *Estuarine Coastal and Shelf Science* 82, 294–300. http://dx.doi.org/10.1016/j.ecss.2009.01.012
- Fredriksson, R., Bergström, U., & Bergström, L. (2010). *Kartläggning av viktiga livsmiljöer för fisk i grunda områden i Kattegatt*. Fiskeriverkets kustlaboratorium. FINFO 2010:4. (In Swedish).
- Gunnartz, U., Lif, M., Lindberg, P., Ljunggren, L., Sandström, A., & Sundblad, G. (2011). *Kartläggning av lekområden för kommersiella fiskarter längs den svenska ostkusten* – *en intervjustudie*. FINFO 2011:3. 42 p. (In Swedish, with English summary).
- HELCOM (1998). Guidelines for Monitoring Phytobenthic Plant and Animal Communities in the Baltic Sea. www.helcom.fi
- HELCOM (2007). Baltic Sea Action Plan. HELCOM Ministerial Meeting: Krakow, Poland, November 17, no. 2007:101.
- HELCOM (2012). Indicator based assessment of coastal fish community status in the Baltic Sea 2005–2009. Baltic Sea Environment Proceeding No. 131.
- HELCOM Guidelines (2015). Guidelines for coastal fish monitoring sampling methods of HELCOM. http://www.helcom.fi/Documents/Action%20areas/ Monitoring%20and%20assessment/Manuals%20and%20Guidelines/ Guidelines%20for%20Coastal%20fish%20Monitoring%20of%20HELCOM.pdf
- Herkül, K. (2014). Modelling the distribution of habitats and key species in Estonian territorial waters. www.kik.ee/sites/default/files/Uuringud/ aruanne pohi modelleerimine-1.pdf
- Jackson, J.B.C., Kirby, M.X., Berger, W.H., Bjorndal, K.A., Botsford, L.W., Bourque, B.J., Bradbury, R., Cooke, R., Erlandson, J., Estes, J.A., Hughes, T.P., Kidwell, S., Lange, C.B., Lenihan, H.S., Pandolfi, J.M., Peterson, C.H., Steneck, R.S., Tegner, M.J., & Warner, R. (2001). Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293, 629–638. http://dx.doi.org/10.1126/science.1059199
- Jokinen, H., Wennhage, H., Lappalainen, A., Ådjers, K., Rask, M., & Norkko, A. (2015). Decline of flounder (*Platichthys flesus* (L.)) at the margin of the species' distribution range. *Journal of Sea Research* 105, 1–9. http://dx.doi.org/10.1016/ j.seares.2015.08.001
- Jokinen, H., Wennhage, H., Ollus, V., Aro, E., & Norkko, A. (2016). Juvenile flatfish in the northern Baltic Sea long-term decline and potential links to habitat characteristics. *Journal of Sea Research* 107, 67–75. http://dx.doi.org/10.1016/j.seares.2015.06.002
- Kallasvuo, M. (2010). Coastal environmental gradients Key to reproduction habitat mapping of freshwater fish in the Baltic Sea. PhD-thesis, Faculty of Biological and Environmental Sciences, University of Helsinki. http://urn.fi/URN:ISBN:978-952-10-6392-3.
- Katsanevakis, S., Wallentinus, I., Zenetos, A., Leppäkoski, E., Çinar, M.E., Oztürk, B., Grabowski, M., Golani, D., & Cardoso, A.C. (2014). Impacts of invasive alien marine species on ecosystem services and biodiversity: a pan-European review. *Aquatic Invasions* 9, 391–423. http://dx.doi.org/10.3391/ai.2014.9.4.01
- Knospina, E. & Putnis, I. (2014). Round Goby Increasingly Guest in the Latvian Coastal Area (Apaļais jūrasgrundulis – aizvien biežāks viesis Latvijas piekrastē). *Latvijas zivsaimniecība* 2014, Rīga, pp. 46–51 (in Latvian).

- Korpinen, S., Jormalainen, V., & Honkanen, T. (2007). Bottom–up and cascading top– down control of macroalgae along a depth gradient. *Journal of Experimental Marine Biology and Ecology* 343, 52–63. http://dx.doi.org/10.1016/j.jembe.2006.11.012
- Kristensen, L.D., Støttrup, J.G., Andersen, S. K., & Degel, H. (2014). Registrering af fangster i de danske kystområder med standardredskaber. Nøglefiskerrapport 2011– 2013. Institut for Akvatiske Ressourcer, Danmarks Tekniske Universitet, DTU Aqua-rapport nr. 286-2014. 100 pp. (In Danish).
- Lotze, H.K. (2004). Repetitive history of resource depletion and mismanagement: the need for a shift in perspective. *Marine Ecology Progress Series* 274, 269–303.
- Lotze, H.K., Lenihan, H.S., Bourque, B.J., Bradbury, R.H, Cooke, R.G., Kay, M.C., Kidwell, S.M., Kirby, M.X., Peterson, C.H., &Jackson, J.B.C. (2006). Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science* 312, 1806–1809. http://dx.doi.org/10.1126/science.1128035
- Martin, G., Möller, T., Kotta, J., Daunys, D., Jermakovs, V., Bucas, M., Siaulys, A., Saskov, A., & Aigars, J. (2010). *Benthic marine habitats of Eastern Baltic Sea*. Tartu Ülikool, Eesti Mereinstituut.
- Martinsson, J. (2011). *Ecology of juvenile turbot and flounder in the central Baltic Sea. Implications for recruitment.* PhD-thesis, Department of Systems Ecology, Stockholm University.
- Moksnes, P.O., Gullström, M., Tryman, K., & Baden, S. (2008). Trophic cascades in a temperate seagrass community. *Oikos* 117, 763–777. http://dx.doi.org/10.1111/j.0030-1299.2008.16521.x
- Möllmann, C., Lindegren, M., Blenckner, T., Bergström, L., Casini, M., Diekmann, R., Flinkman, J., Müller-Karulis, B., Neuenfeldt, S., Schmidt, J.O., Tomczak, M., Voss, R., & Gårdmark, A. (2014). Implementing ecosystem-based fisheries management: from single-species to integrated ecosystem assessment and advice for Baltic Sea fish stocks. *ICES Journal of Marine Science* 71, 1187–1197. http://dx.doi.org/ 10.1093/icesjms/fst123
- Murray, G., Neis, B., Palmer, C.T., & Schneider, D.C. (2008a). Mapping cod: fisheries science, fish harvesters' ecological knowledge and cod migrations in the northern Gulf of St. Lawrence. *Human Ecology* 36, 581–598. http://dx.doi.org/10.1007/s10745-008-9178-1
- Murray, G., Neis, B., Schneider, D.C., Ings, D., Gosse, K., Whalen, J., & Palmer, C.T. (2008b). Local ecological knowledge in the historical reconstruction of marine socioenvironmental systems: Methods, procedures and challenges. In: Neis, B., Lutz, J. (Eds.), *Making and Moving Knowledge: Interdisciplinary and Community-based Research for a World on the Edge*, McGill-Queen's University Press, pp. 100–120.
- Neis, B., Schneider, D.C., Felt, L.F., Haedrich, R.L., Hutchings, J.A., & Fischer, J. (1999). Northern cod stock assessment: What can be learned from interviewing resource users? *Canadian Journal of Fisheries and Aquatic Sciences* 56, 1944–1963. http://dx.doi.org/10.1139/f99-115
- Newcombe, E.M., & Taylor, R.B. (2010). Trophic cascade in a seaweed-epifauna-fish food chain. *Marine Ecology Progress Series* 408, 161–167. http://dx.doi.org/ 10.3354/meps08589
- Nissling, A., Jacobsson, M., & Hallberg, N. (2007). Feeding ecology of juvenile turbot, *Scophthalmus maximus* and flounder *Pleuronectes flesus* at Gotland, Central Baltic Sea. *Journal of Fish Biology* 70, 1877–1897. http://dx.doi.org/10.1111/j.1095-8649.2007.01463.x

- Nohrén, E., Pihl, L., & Wennhage, H. (2009). Spatial patterns in community structure of motile epibenthic fauna in coastal habitats along the Skagerrak–Baltic salinity gradient. *Estuarine Coastal and Shelf Science* 84, 1–10. http://dx.doi.org/10.1016/j.ecss.2009.05.012
- Norderhaug, K.M., Christie, H., Fosså, J.H., & Fredriksen, S. (2005). Fish-macrofauna interactions in a kelp (*Laminaria hyperborea*) forest. *Journal of the Marine Biological Association of the UK* 83, 1279–1286. http://dx.doi.org/10.1017/S0025315405012439
- Nyström Sandman, A., Didrikas, T., Enhus, C., Florén, K., Isæus, M., Nordemar, I., Nikolopoulos, A., Sundblad, G., Svanberg, K., & Wijkmark, N. (2013a). *Marin Modellering i Södermanlands län, AquaBiota Report 2013:09*. (In Swedish).
- Nyström Sandman, A., Didrikas, T., Enhus, C., Florén, K., Isæus, M., Nordemar, I., Nikolopoulos, A., Sundblad, G., Svanberg, K., & Wijkmark, N. (2013b). *Marin Modellering i Stockholms län, AquaBiota Report 2013:10*. (In Swedish).
- Öhnstedt, E. (2015). Modellering av tånglakens, *Zoarces viviparus*, habitat i Östersjön. Modelling of the eelpout's, *Zoarces viviparus*, habitat in the Baltic Sea. Independent project in Biology 15 HEC, SLU, http://stud.epsilon.slu.se. (In Swedish).
- Ojaveer, E., Arula, T., Lankov, A., & Shpilev, H. (2011). Impact of environmental deviations on the larval and year-class abundances in the spring spawning herring (*Clupea harengus membras* L.) of the Gulf of Riga (Baltic Sea) in 1947–2004. *Fisheries Research* 107, 159–168. http://dx.doi.org/10.1016/j.fishres.2010.11.001
- Ojaveer, H. & Kotta, J. (2015). Ecosystem impacts of the widespread non-indigenous species in the Baltic Sea: literature survey evidences major limitations in knowledge. *Hydrobiologia* 750, 171–185. http://dx.doi.org/10.1007/s10750-014-2080-5
- Olsson, J., Bergström, L. & Gårdmark, A. (2012). Abiotic drivers of coastal fish community change during four decades in the Baltic Sea. *ICES Journal of Marine Science* 69, 961–970. http://dx.doi.org/10.1093/icesjms/fss072
- Olsson, J., Bergström, L. & Gårdmark, A. (2013a). Top-down regulation, climate and multi-decadal changes in coastal zoobenthos communities in two Baltic Sea areas. *PLoS One*, e64767. http://dx.doi.org/10.1371/journal.pone.0064767
- Olsson, J., Kaljuste, M., Heimbrand, Y., Odelström, A., & Bergström, U. (2013b). *Otolitkemisk analys av abborre och gädda i Blekinge och östra Skåne*. Rapport till länsstyrelsen i Blekinge.
- Orth, R.J., Carruthers, T.J., Dennison, W.C., Duarte, C.M., Fourqurean, J.W., Heck, K.L., Hughes, A.R., Kendrick, G.A., Kenworthy, W.J., Olyarnik, S., Short, F.T., Waycott, M., & Williams, S.L. (2006). A global crisis for seagrass ecosystems. *Bioscience* 56, 987– 996. http://dx.doi.org/10.1641/0006-3568(2006)56[987:AGCFSE]2.0.C0;2
- Östman, Ö., Eklöf, J., Eriksson, B.K., Olsson, J., Moksnes, P.-O., & Bergström, U. (2016). Top-down control as important as nutrient enrichment for eutrophication effects in North Atlantic coastal ecosystems. *Journal of Applied Ecology* 53:1138-1147. http://dx.doi.org/10.1111/1365-2664.12654
- Pikitch, E.K., Santora, C., Babcock, E.A., Bakun, A., Bonfil, R., Conover, D.O., Dayton, P.K., Doukakis, P., Fluharty, D., Heneman, B., Houde, E.D., Link, J., Livingston, P.A., Mangel, M., McAllister, M.K., Pope, J., & Sainsbury, K.J. (2004). Ecosystem-based fishery management. *Science* 305, 346–347. http://dx.doi.org/10.1126/ science.1098222

Rangeley, R.W., & Kramer, D.L. (1995). Tidal effects on habitat selection and aggregation by juvenile Pollock *Pollachius virens* in the rocky intertidal zone. *Marine Ecology Progress Series* 126, 19–29. http://dx.doi.org/10.3354/ meps126019

Rohtla, M., Vetemaa, M., Urtson, K., & Soesoo, A. (2012). Early life migration patterns of Baltic Sea pike *Esox lucius. Journal of Fish Biology* 80, 886–893. http://dx.doi.org/10.1111/j.1095-8649.2012.03226.x

Rohtla, M., Vetemaa, M., Taal, I., Svirgsden, R., Urtson, K., Saks, L., Verliin, A., Kesler, M., & Saat, T., (2014). Life history of anadromous burbot (*Lota lota*, Linneaus) in the brackish Baltic Sea inferred from otolith microchemistry. *Ecology of Freshwater Fish* 23, 141–148. http://dx.doi.org/10.1111/eff.12057

Rohtla, M., Svirgsden, R., Taal, I., Saks, L., Eschbaum, R., & Vetemaa, M. (2015). Lifehistory characteristics of ide *Leuciscus idus* in the Eastern Baltic Sea. *Fisheries Management and Ecology* 22, 239–248. http://dx.doi.org/10.1111/fme.12120

Rousi, H., Laine, A.O., Peltonen, H., Kangas, P., Andersin, A.B., Rissanen, J., Sandberg-Kilpi, E., & Bonsdorff, E. (2013). Long-term changes in coastal zoobenthos in the northern Baltic Sea: the role of abiotic environmental factors. *ICES Journal of Marine Science* 70, 440–451. http://dx.doi.org/10.1093/icesjms/fss197

Saat, T. (2015). Kalakoelmute seisund ning koelmualade melioreerimise lähteülesannete koostamine. (In Estonian) https://www.google.ee/ webhp?sourceid=chrome-instant&ion=1&espv=2&ie=UTF-8#q=r%C3%A4ime%20koelmualade%20modelleerimine

Sandström, A., Eriksson, B.K., Karås, P., Isæus, M., & Schreiber, H. (2005). Boating and navigation activities influence the recruitment of fish in a Baltic Sea archipelago area. *Ambio* 34, 125–130. http://dx.doi.org/10.1579/0044-7447-34.2.125

Šaškov, A., Šiaulys, A., Bučas, M., & Daunys, D. (2014). Baltic herring (*Clupea harengus membras*) spawning grounds on the Lithuanian coast: current status and shaping factors. *Oceanologia* 56, 789–804. http://dx.doi.org/10.5697/oc.56-4.789

Seitz, R.D., Wennhage, H., Bergström, U., Lipcius, R.N., & Ysebaert, T. (2014). Ecological value of coastal habitats for commercially and ecologically important species. *ICES Journal of Marine Science* 71, 648–665. http://dx.doi.org/10.1093/icesjms/fst152

Sieben, K., Ljunggren, L., Bergström, U., & Eriksson, B.K. (2011). A mesopredator release of stickleback promotes recruitment of macroalgae in the Baltic Sea. *Journal of Experimental Marine Biology and Ecology* 397, 79–84. http://dx.doi.org/10.1016/j.jembe.2010.11.020

Snickars, M., Sandström, A., Lappalainen, A., & Mattila, J. (2007). Evaluation of low impact pressure waves as a quantitative sampling method for small fish in shallow water. *Journal of Experimental Marine Biology and Ecology* 343, 138–147. http://dx.doi.org/10.1016/j.jembe.2006.12.008

Snickars, M., Sundblad, G., Sandström, A., Ljunggren, L., Bergström, U., Johansson, G., & Mattila, J. (2010). Habitat selectivity of substrate-spawning fish: modelling requirements for the Eurasian perch *Perca fluviatilis. Marine Ecology Progress Series* 398, 235–243. http://dx.doi.org/10.3354/meps08313

Snickars, M., Weigel, B., & Bonsdorff, E. (2015). Impact of eutrophication and climate change on fish and zoobenthos in coastal waters of the Baltic Sea. *Marine Biology* 162, 141–151. http://dx.doi.org/10.1007/s00227-014-2579-3

- Stål, J., Pihl, L., & Wennhage, H. (2007). Food utilisation by coastal fish assemblages in rocky and soft bottoms on the Swedish west coast: Inference for identification of essential fish habitats. *Estuarine Coastal and Shelf Science* 71, 593–607. http://dx.doi.org/10.1016/j.ecss.2006.09.008
- Sundblad, G., & Bergström, U. (2014). Shoreline development and degradation of coastal fish reproduction habitats. *Ambio* 43, 1020–1028. http://dx.doi.org/10.1007/s13280-014-0522-y
- Sundblad, G., Bergström, U., & Sandström, A. (2011). Ecological coherence of marine protected area networks: a spatial assessment using species distribution models. *Journal of Applied Ecology* 48, 112–120. http://dx.doi.org/10.1111/j.1365-2664.2010.01892.x
- Sundblad, G., Bergström, U., Sandström, A., & Eklöv, P. (2014). Nursery habitat availability limits adult stock sizes of predatory coastal fish. *ICES Journal of Marine Science* 71, 672–680. http://dx.doi.org/10.1093/icesjms/fst056
- Svensson, C.J., Baden, S., Moksnes, P.O., & Åberg, P. (2012). Temporal mismatches in predator-herbivore abundance control algal blooms in nutrient-enriched seagrass ecosystems. *Marine Ecology Progress Series* 471, 61–71. http://dx.doi.org/10.3354/meps10014
- Taal, I., Kesler, M., Saks, L., Rohtla, M., Verliin, A., Svirgsden, R., Jürgens, K., Vetemaa, M., & Saat, T. (2014). Evidence for an autumn downstream migration of Atlantic salmon *Salmo salar* (Linnaeus) and brown trout *Salmo trutta* (Linnaeus) parr to the Baltic Sea. *Helgoland Marine Research* 68, 373–377. http://dx.doi.org/10.1007/s10152-014-0395-2
- Turner, S.J., Thrush, S.F., Hewitt, J.E., Cummings, V.J., & Funnell, G. (1999). Fishing impacts and the degradation or loss of habitat structure. *Fisheries Management and Ecology* 6, 401–420. http://dx.doi.org/10.1046/j.1365-2400.1999.00167.x
- Ubl, C. & Dorow, M. (2015). A novel enclosure approach to assessing yellow eel (*Anguilla anguilla*) density in non-tidal coastal waters. *Fisheries Research* 161, 57–63. http://dx.doi.org/10.1016/j.fishres.2014.06.009
- Ustups, D., Müller-Karulis, B., Bergström, U., Makarchouk, A., & Sics, I. (2013). The influence of environmental conditions on early life stages of flounder (*Platichthys flesus*) in the central Baltic Sea. *J Sea Res* 75, 77–84. http://dx.doi.org/10.1016/ j.seares.2012.05.001
- Vanhatalo, J., Veneranta, L., & Hudd, R. (2012). Species distribution modeling with Gaussian processes: A case study with the youngest stages of sea spawning whitefish (*Coregonus lavaretus* L. sl) larvae. *Ecological Modelling* 228, 49–58. http://dx.doi.org/10.1016/j.ecolmodel.2011.12.025
- Veneranta, L., Urho, L., Lappalainen, A., & Kallasvuo, M. (2011). Turbidity characterizes the reproduction areas of pikeperch (*Sander lucioperca* (L.)) in the northern Baltic Sea. *Estuarine Coastal and Shelf Science* 95, 199–206. http://dx.doi.org/10.1016/j.ecss.2011.08.032
- Vetemaa, M., Eschbaum, R., Verliin, A., Albert, A., Eero, M., Lillemägi, R., Pihlak, M., & Saat, T. (2006). Annual and seasonal dynamics of fish in the brackish-water Matsalu Bay, Estonia. *Ecology of Freshwater Fish* 15, 211–220. http://dx.doi.org/10.1111/j.1600-0633.2006.00134.x
- Warnar, T., Huwer, B., Vinther, M., Egekvist, J., Sparrevohn, C.R., Kirkegaard, E., Dolmer, P., Munk, P., & Sørensen, T.K. (2012). *Fiskebestandenes struktur. Fagligt baggrundsnotat til den danske implementering af EU's havstrategidirektiv.* Institut for Akvatiske Ressourcer, Danmarks Tekniske Universitet. (In Danish).

Weigel, B., Andersson, H.C., Meier, H.M., Blenckner, T., Snickars, M., & Bonsdorff, E., (2015). Long-term progression and drivers of coastal zoobenthos in a changing system. *Marine Ecology Progress Series* 528, 141–159. http://dx.doi.org/10.3354/ meps11279

Wijkmark, N. (ed.) (2014). Field, laboratory and experimental work within the MAR-MONI project – report on survey results and obtained data. pp. 104. http://www.aquabiota.se/wp-content/uploads/A3_REPORT_Survey_Results.pdf

Willebrand, S. (2015). Habitat preference of *Neogobius melanostomus* in the Baltic Sea. Independent project in Biology 15 HEC, SLU, http://stud.epsilon.slu.se

http://paikkatieto.ymparisto.fi/velmu [page visited 18.8.2016]

http://pump.regio.ee/kalandus/public/ [page visited 18.8.2016]

Svensk sammanfattning

Många fiskarter i Östersjön är beroende av grunda och skyddade kusthabitat för lek, uppväxt, födosök och vid migration. Dessa essentiella fiskhabitats (EFH) roll för utvecklingen av fiskbestånd och fiskproduktion har trots detta fått relativt lite uppmärksamhet, fastän man inom vetenskapen är överens om deras kritiska betydelse. Man vet också ganska lite om de viktigaste hoten och påverkanstrycken mot EFH, samt deras bevarandestatus i olika Östersjöländer. Eftersom EFH ofta förekommer i de delar av kustzonen som värderas högt av människan, bör dessa kunskapsluckor fyllas. Följaktligen finns det ett akut behov av att mera grundligt fokusera på betydelsen, kartläggningen, övervakningen och skyddet av EFH och också på påverkanstryck och mekanismer bakom de förändringar vi kan observera i habitatens tillstånd. Detta för att kunna förutspå och minska framtida effekter av miljöförändringar i habitaten och kunna införa långsiktiga och anpassade förvaltningsplaner för EFH.

Målsättningarna med detta projekt var 1) att organisera en workshop med inbjudna experter kring Östersjön för att diskutera betydelsen och skyddet av samt hoten mot kustnära EFH (inklusive en sammanställning av metoder som används för kartläggning och övervakning av dessa habitat i olika länder), samt 2) på basen av resultaten från denna workshop skriva en artikel för en internationell vetenskaplig tidskrift kring det nuvarande kunskapsläget inom området i Östersjön, inklusive kunskapsluckor och framtida forskningsbehov. I denna rapport presenterar vi resultaten från detta projekt med specifikt fokus på utfallet av workshopen.

Från workshopen, som ordnades 2–4 juni 2015 i Öregrund i Sverige, kan man dra slutsatsen att det bara finns ett fåtal kvantitativa undersökningar från Östersjön gällande betydelsen av EFH för fiskbestånd. Bevisen är i sin tur rätt komplexa och ger inte nödvändigtvis enkla svar. Inte desto mindre finns det för några arter av kustfisk indirekta bevis och tillräcklig mängd data för att genomföra fortsatta kvantitativa analyser. Mera bevis om betydelsen av EFH för fiskproduktion kan möjligen erhållas genom fortsatt användning av rumsliga och tidsmässiga analyser, olika modelleringstekniker som fokuserar på fiskarnas olika livsstadier (t.ex. stadiestrukturerade modeller), samt otolitkemiska tekniker. Baserat på den information som finns att tillgå är EFH väldigt viktiga och värdefulla för tillhandahållandet av friska fisksamhällen och för produktion av fisk. Troligen har betydelsen av EFH underskattats och framtida undersökningar skulle kunna bidra med att klarlägga deras ekologiska betydelse.

Gällande olika aspekter kring övervakning och kartläggning av EFH i Östersjön verkar det finnas en hel del data tillgängligt. Olika provtagningsmetoder används för en stor mängd av fiskarter och livsstadier (från ägg och juveniler till vuxen fisk) och habitat (från kustnära områden till det öppna havet). Nyttjandet av dessa data som underlag för habitatkartor har länge varit relativt lågt. Situationen håller ändå nu på att förändras i många länder genom nyligen startade nationella och regionala program för inventering och kartläggning av undervattensmiljöer. Viktigt i detta sammanhang är att man i kartläggningen även bör fokusera på huvudsakliga hot, påverkanstryck och bakgrundsförhållanden för fisken och dess miljö. Detta kunde säkerställa en optimal tillgång till och användning av information som är väsentlig för en effektiv förvaltning och för en förbättrad marin planering.

Hoten mot, och bevarandestatusen för, EFH i Östersjön kräver brådskande och mångsidiga förvaltningslösningar. Eutrofiering, klimatförändring, kustexploatering, främmande arter och fiske verkar utgöra de främsta hoten för habitaten. Bland dessa hot är de fysiska påverkanstrycken (omfattande t.ex. marin trafik och dess infrastruktur såsom muddring), fysisk exploatering av strandmiljöer och trålfiske kanske de som är lättast att hantera förvaltningsmässigt. Dessa aktiviteter påverkar fiskhabitat både direkt och indirekt, och utgör i regel störst hot mot fiskens reproduktion och tidiga uppväxt (yngelstadier). Bevarandestatusen för EFH är i allmänhet låg, vilket huvudsakligen är en följd av att fiskeriförvaltningen och miljöförvaltningen historiskt sett har varit åtskilda i Östersjöregionen. Flera studier har dock visat att fördelar kan uppnås genom en integrerad förvaltning av fisken och dess habitat. Genom att föra samman fiskeriförvaltning med miljöförvaltning (t.ex. habitatskydd) får EFH en central roll inom förvaltningen samtidigt som också vetenskapliga intressen för relevanta frågeställningar för forskningen väcks.

Resultaten från detta projekt visar att 1) det finns tillgängliga data för kvantitativa analyser som kan stöda rollen av EFH för fiskproduktion, 2) det finns en potential för att initiera, utveckla och synkronisera framtida övervakning och kartläggning av habitaten, samt 3) att det finns en ökad medvetenhet kring skyddet av och ett hållbart nyttjande av dessa grunda kustnära system. Resultaten från detta projekt kan fungera som ett underlag för ett förbättrat samarbete mellan Östersjöländerna inom detta fält, vilket i det långa loppet kunde resultera i harmoniserade övervaknings- och kartläggningsmetoder för EFH i Östersjön samt stärkt förvaltning och skydd. Arbetet ger också ett viktigt bidrag till utvecklingen av indikatorer för att bedöma tillståndet av EFH och för genomförande av internationella överenskommelser och förordningar såsom HELCOMs Aktionsplan för Östersjön (Baltic Sea Action Plan, BSAP), Habitatdirektivet (HD) och Havsmiljödirektivet (Marine Strategy Framework Directive, MSFD).

Appendix 1 – Participants list

Table 2: Participants

Name	Last name	Organization	Country	E-mail address
Meri	Kallasvuo	Natural Resources Institute	Finland	Meri.Kallasvuo@luke.fi
Antti	Lappalainen	Natural Resources Institute	Finland	Antti.Lappalainen@luke.fi
Henri	Jokinen	University of Helsinki	Finland	henri.jokinen@helsinki.fi
Martin	Snickars	Åbo Akademi University	Finland	martin.snickars@abo.fi
Göran	Sundblad	AquaBiota Water Research	Sweden	goran.sundblad@aquabiota.se
Helmut	Winkler	University of Rostock	Germany	helmut.winkler@uni-rostock.de
Peter	Möller	Fisch und Umwelt, Mecklenburg-Vor- pommern	Germany	p.moeller@fischumwelt.de
Josianne	Støttrup	DTU Aqua	Denmark	jgs@aqua.dtu.dk
Elliot John	Brown	DTU Aqua	Denmark	elbr@aqua.dtu.dk
Linas	Lozys	Nature Research Centre	Lithuania	lozys@ekoi.lt
Justas	Dainys	Nature Research Centre	Lithuania	justas@akvaklubas.lt
Laura	Briekmane	Institute of Food Safety, Animal Health and Environment	Latvia	Laura.Briekmane@bior.lv
Didzis	Ustups	Institute of Food Safety, Animal Health and Environment	Latvia	Didzis.Ustups@bior.lv
Adam	Wozniczka	National Marine Fisheries Research Institute	Poland	awozniczka@mir.gdynia.poland
Zuzanna	Celmer	National Marine Fisheries Research Institute	Poland	zcelmer@mir.gdynia.poland
Timo	Arula	Estonian Marine Institute, University of Tartu	Estonia	timo.arula@ut.ee
Aare	Verliin	Estonian Marine Institute, University of Tartu	Estonia	aare.verliin@ut.ee
Lauri	Saks	Estonian Marine Institute, University of Tartu	Estonia	slauris@ut.ee

Name	Last name	Organization	Country	E-mail address
Imre	Taal	Estonian Marine Institute, University of Tartu	Estonia	imre.taal@ut.ee
Markus	Vetemaa	Estonian Marine Institute, University of Tartu	Estonia	markus.vetemaa@ut.ee
Mehis	Rohtla	Estonian Marine Institute, University of Tartu	Estonia	mehis.rohtla@ut.ee
Alessandro	Orio	Swedish University of Agricultural Sciences, SLU-Aqua	Sweden	alessandro.orio@slu.se
Patrik	Kraufvelin	Swedish University of Agricultural Sci- ences, SLU-Aqua	Sweden	patrik.kraufvelin@slu.se
Ann-Britt	Florin	Swedish University of Agricultural Sci- ences, SLU-Aqua	Sweden	ann-britt.florin@slu.se
Ulf	Bergström	Swedish University of Agricultural Sci- ences, SLU-Aqua	Sweden	ulf.bergstrom@slu.se
Zeynep	Pekcan-Hekim	Swedish University of Agricultural Sciences, SLU-Aqua	Sweden	zeynep.pekcan.hekim@slu.se
Annukka	Lehikoinen	Swedish University of Agricultural Sci- ences, SLU-Aqua	Sweden	annukka.lehikoinen@slu.se
Jens	Olsson	Swedish University of Agricultural Sciences, SLU-Aqua	Sweden	jens.olsson@slu.se
Johanna	Mattila	Swedish University of Agricultural Sci- ences, SLU-Aqua	Sweden	johanna.mattila@slu.se
Petra	Kääriä	HELCOM Secretariat	Finland	Petra.Kaaria@helcom.fi

Appendix 2 – Schedule of the workshop

Essential Coastal Habitats for Fish Workshop (2nd-4th June, Öregrund).

2nd June Tuesday Day 1		
13:00-14:00	Lunch at Strandhotellet	
14:00	Start – Welcome	
	Introduction of the Institute of Coastal Research and SLU – Zeynep Pekcan-Hekim	
	Introducing the workshop program and themes – Jens Olsson	
Theme 1: The role of essential coastal habitats for fish – availability of quantitative evidence showing the importance of coastal habitats for fish stocks		
14:30	Göran Sundblad – Habitat population size relationships	
14:45	Josianne Støttrup – Identifying and quantifying essential fish habitats	
15:00	Henri Jokinen – Essential habitats and (unresolved) links to fish stocks on the Finnish coast: examples from flounder studies	
15:15	Adam Woźniczka – The role of Baltic lagoon on forming the fish communities in coastal wa- ters. Szczecin Lagoon – Pomeranian Bay case studies	
15:30	Coffee Break (30 mins)	
16:00	Didzis Ustups – Habitat use by early life stages of flounder in the Central Baltic Sea	
16:15	Mehis Rohtla – Spawning habitat preference of Väinameri pike, ide and burbot	
16:30	Timo Arula – Clupeid spawning ground surveys in the NE Baltic Sea	
16:45	End of day – wrapping up day1	
18:30	Boat Trip – meeting at the harbor see map	

Table 4: Schedule of the workshop, Day 2

3rd June Wednesday Day 2		
Theme 2: How are essential coastal habitats for fish mapped and monitored? – methods, available maps, are there data for maps?		
9:00	Introduction to day 2	
9:15	Meri Kallasvuo – Using high-resolution species distribution modelling to produce reproduction habitat maps of coastal fish to support marine spatial planning	
9:30–12	Working group on Theme 2 methods by country	
12–13	Lunch at Strandhotellet	
Theme 3: Conse	ervation and threats on essential coastal habitats for fish	
13:00	Patrik Kraufvelin – Physical threats to essential coastal habitats for fish	
13:15	Ulf Bergström – Essential fish habitats in management	
13:30	Antti Lappalainen – Conservation and threats on essential coastal habitats for fish – Finland	
13:45	Elliot John Brown – Conservation and threats on essential coastal habitats for fish: The need for protection of essential fish habitats	
14:00	Lina Lozys and Justas Dainys – Essential coastal habitats for fish in Lithuania: threats and conservation	
14:15	Coffee Break (30 mins)	
14:45	Martin Snickars – Vulnerable Nursery Habitats – Coastal Lagoons	
15:00	Peter Möller – Bottom habitat measures within the coastal areas of the State of Mecklenburg- Western Pomerania	
15:15	Lauri Saks – Coastal marine habitats and their conservation status in Estonia	
15:30	Zuzanna Celmer – Coastal fish habitats the Puck Bay case	
15:45	Helmut Winkler – Coastal fish habitats in the Pomeranian Bay	
16:00-17:00	Discussion on the current needs and trends in different countries on the protection of the es- sential coastal habitats for fish	
19:00	Dinner at Societetshuset	

Table 5: Schedule of the workshop, Day 3

4th June Thursday Day 3 (Meeting moves to SLU)		
9:00-12:00	Presentations on <i>Theme 2</i> summing up group work (By country) Continue discussions on <i>Theme 1</i> looking for evidence (Everyone)	
12:00-13:00	Lunch at Strandhotellet	
13:00-15:00	General discussion on the outcomes of the workshop Drafting the review Discussion on potential funding possibilities	

Appendix 3 – Abstracts from presentations

Abstract Jens Olsson:Introduction to the workshop

Fish are important due to many reasons: in economic terms (commercial species); in socio-economical terms (coastal fish species - recreational fisheries); for ecosystem functioning - predatory fish counteract eutrophication symptoms; as indicators for ecosystem status and health; as segments of the MSFD and BSAP, as well as in focus of the CFP. In the Baltic Sea, the occurrence/distribution of fish is structured by many background factors such as: salinity, temperature, predation, pollution, nutrient enrichment, reproduction, fishery and finally, habitats - the central topic of this workshop. With this in mind and for the central purposes of the workshop, the following specific gaps in knowledge may be emphasized: the role of EFH is relatively little studied; the need for characterizing, mapping and monitoring of EFH; there is a lack of information about the most important threats to and their effects on these habitats; the fact that coastal habitats are under national jurisdiction (which potentially is affecting conservation and management issues). The workshop has specifically been organized for the following central target groups: for ourselves, for HELCOM, for ICES and for the scientific community, with the major reason being to search for ways in how to better connect fisheries and environmental management. The basic timeline for the work to be carried out looks as follows: workshop - June 2015; reporting of the outcome of the workshop - November 2015; writing a scientific review - late 2015 – early 2016; possible presentation at HELCOM FISH – 2016; possible common research proposal/application – 2016; preparation of a HEL-COM guideline for mapping and monitoring EFH – 2018.

Abstracts from *Theme 1* – The role of essential coastal habitats for fish –

availability of quantitative evidence showing the importance of coastal habitats for fish stocks

Abstract Göran Sundblad

Habitat population size relationships

Many Baltic Sea fish species are dependent on coastal habitats. A recent review have shown that 44% of all ICES species utilize coastal habitats as spawning, feeding, nursery or migration areas, and these stocks contribute 77% of the commercial landings of ICES-advice species (Seitz et al. 2013). Additionally, by intuition, we know that a limited habitat supply must, by some means, control the size and dynamics of fish populations, but how can this be quantified? For species with ontogenetic habitat shifts, species distribution modelling has emerged as a tool to map lifestage specific habitat requirements. By relating the distribution of nursery habitats for perch and pikeperch to the adult population size in twelve archipelago areas between Sweden and Finland, the importance of nursery habitats could be quantified (Sundblad et al. 2014). Habitat availability explained almost half of the variation in population size, indicating that habitat availability plays a crucial role in limiting adult stock sizes. The relationships were non-linear suggesting that the effect of habitat loss or restoration would be largest in areas with little available habitat. Importantly, a map based spatial approach yields quantitative links, which allows for map based estimates of the regional distribution of large fish, scenario analyses etc. (e.g. Bergström et al. 2013).

It appears that the scientific literature is not abundant with quantitative evidence for the importance of coastal habitats, but some studies are available. These include i) model based approaches (e.g. Halpern *et al.* 2005, Minns *et al.* 1996), ii) the importance of specific habitat types using elemental fingerprinting (e.g. Fodrie and Levin 2008), habitat specific biomass and size distributions (e.g. Mumby *et al.* 2004) and nursery habitat size (Rijnsdorp *et al.* 1992), as well as iii) long term field experiments (Schmitt and Holbrook 2000), perhaps providing the strongest evidence for the importance of specific coastal fish habitat types.

Studies of fish populations appear to have evolved along two distinct paths, where one has centered on the dynamics of populations in relation to their exploitation and another has focused on the static relationships between fish and their habitats. As quantitative evidence for habitat limitation is accumulating, from different areas, species and with different methods for quantification, there is now an unprecedented possibility, as well as need, to integrate habitat in fisheries management and nature conservation. Especially in the light of efforts towards an ecosystem based approach to (fisheries) management.

References

- Bergström, U., G. Sundblad, A.-L. Downie, M. Snickars, C. Boström, & M. Lindegarth. (2013). Evaluating eutrophication management scenarios in the Baltic Sea using species distribution modelling. *Journal of Applied Ecology 50*:680–690. http://dx.doi.org/10.1111/1365-2664.12083
- Fodrie, F. J. & L. A. Levin (2008). Linking juvenile habitat utilization to population dynamics of California halibut. *Limnology and oceanography* 53:799–812. http://dx.doi.org/10.4319/lo.2008.53.2.0799
- Halpern, B. S., S. D. Gaines, & R. R. Warner (2005). Habitat size, recruitment, and longevity as factors limiting populations size in stage-structured species. *The American Naturalist* 165:82–94. http://dx.doi.org/10.1086/426672
- Mumby, P. J., A. J. Edwards, J. E. Arias-González, K. C. Lindeman, P. G. Blackwell, A. Gall, M. I. Gorczynska, A. R. Harborne, C. L. Pescod, H. Renken, C. C. C. Wabnitz, & G. Llewellyn (2004). *Mangroves enhance the biomass of coral reef fish communities in the Carribean. Nature* 427:533–536. http://dx.doi.org/10.1038/nature02286
- Minns, C. K., R. G. Randall, J. E. Moore, & V. W. Cairns (1996). A model simulating the impact of habitat supply limits on northern pike, Esox lucius, in Hamilton Harbour, Lake Ontario. *Canadian Journal of Fisheries and Aquatic Sciences* 53:20–34. http://dx.doi.org/10.1139/f95-258
- Rijnsdorp, A. D., F. A. Van Beek, S. Flatman, R. M. Millner, J. D. Riley, M. Giret, & R. De Clerck. 1992. Recruitment of sole stocks, Solea solea (L.), in the Northeast Atlantic. *Netherlands Journal of Sea Research 29*:173–192. http://dx.doi.org/10.1016/0077-7579(92)90018-A
- Schmitt, R. J. & S. J. Holbrook (2000). Habitat-limited recruitment of coral reef damselfish. *Ecology 81*:3479–3494. http://dx.doi.org/10.1890/ 0012-9658(2000)081[3479:HLROCR]2.0.C0;2
- Seitz, R. D., H. Wennhage, U. Bergström, R. N. Lipcius & T. Ysebaert (2013). Ecological value of coastal habitats for commercially and ecologically important species. ICES *Journal of Marine Science* 71:648–665. http://dx.doi.org/10.1093/icesjms/fst152
- Sundblad, G., U. Bergström, A. Sandström & P. Eklöv (2014). Nursery habitat availability limits adult stock sizes of predatory coastal fish. *ICES Journal of Marine Science* 71:672–680. http://dx.doi.org/10.1093/icesjms/fst056

Abstract: Støttrup, J.G., Sørensen, T.K., Egekvist, J., Vinther, M., Brown, J.E., Dinesen, G.E. and Hansen, F.I.

How can we quantify the importance of coastal habitats for fish stocks?

Essential fish habitats include nursery, feeding and spawning areas and migratory routes. The latter three habitats are targeted by fishermen and may be important for fishers because of high catches or value per fishery effort in these areas. Therefore some habitats are important both for fish stocks and targeted by fisheries which creates challenges for their sustainable management. Mapping these habits is essential for management but can be done at different levels. Potential fish habitats may be mapped to indicate where one may expect specific life stages of specific fish to be distributed. Realized habitat is where fish species are monitored as presence or abundance, but effective habitats are those areas with a relatively high contribution to a fish population in comparison to adjacent areas.

The Sound between Denmark and Sweden is a transitional sea between the Baltic and the North Sea with multiple uses such as commercial gillnet fishing, angling and sports diving as well as shipping, the wind energy sector and marine aggregate extraction. Conflicts arise e.g. between the fishing sector and sectors impacting the sea floor and its habitats. A project was therefore carried out in 2014–2015 in the Danish part of the Sound with the aim to collect information from all existing sources. Quantitative information from fishing surveys was lacking and the study relied on general descriptive information and interviews with small-scale commercial gillnetters and anglers to map the habitats of ecological importance for selected fish species of commercial value. The information and any data available was collated in GIS layers and integrated in map overlay using also GIS layers with depth, sediment or other environmental information. Habitat maps for seven commercially important fish species were produced and when compiled provided visualization of hotspot fishery/fish habitat areas, which would be most likely to create conflict between fishery and marine aggregate extraction.

Abstract: Henri Jokinen

Essential habitats and (unresolved) links to fish stocks on the Finnish coast: examples from flounder studies

Essential Coastal Habitats for Fish (EFH) are defined as environments necessary for any life-stage of a species. The importance of the different EFHs can be assessed as the effects of changes in the quantity and/or quality of these habitats on populations or stocks in time or space. The Finnish coast is characterized by a high diversity of habitats. Hence, spatial overlap and mixing between different EFHs is probable, making it difficult to distinguish their effects, and to establish their relative roles for fish stocks. To date only a few studies have tried to link EFHs and fish stocks in the coastal northern Baltic Sea. Shallow littoral areas provide generally good conditions for growth, survival and recruitment of juveniles to the adult population, if the nursery function of these areas is undisturbed, then constituting important coastal EFHs for many species. Flounder on the Finnish coast are facing multiple pressures due to on-going large-scale ecosystem changes linked to e.g. eutrophication and climate change. As a consequence of coastal eutrophication, vegetation coverage and filamentous algae have generally increased in shallow areas. Using available fishery-independent data on adult flounders as well as historical and present-state data on juvenile flatfish in nursery areas, from the Finnish coast, we demonstrate a change over time for both groups. We also examined the present occurrence of juvenile flatfish (flounder and turbot) in relation to vegetation/algae among other environmental factors in shallow juvenile habitats. Based on the available data it seems that both juveniles and adults on the Finnish coast have decline over time, during a period between the 80s and now, but relating these events is difficult because of the large time gap in juvenile data.

Due to restrictions of the data, it was not possible to assess changes over time in juvenile habitat quality or to establish that the decline in juveniles was caused by these changes during the post-settlement stage. The present low occurrence and frequent absence of juveniles in known nurseries further hampered the assessment of juvenile habitat quality on a spatial scale, despite some evidence for association of juveniles to less vegetated areas and for absence of juveniles in the most sheltered and turbid sites. In conclusion, it seems difficult to find quantitative evidence on the role of the juvenile EFHs for the adult flounder stock on the Finnish coast.

Abstract: Didzis Ustups, Ann-Britt Florin, Ulf Bergström and colleagues from BIOR

Habitat use by early life stages of flounder in the Central Baltic Sea Flounder (*Platichthys flesus*) is a temperate marine fish that is well adapted to the brackish waters of the Baltic Sea. There are two sympatric flounder populations in the Baltic Sea, pelagic and demersal spawners, which differ in their spawning habitat and egg characteristics. In the present study, early life stages (eggs, larvae, juveniles) of flounder of the central Baltic Sea were studied. We examined whether variations in hydrological regime can explain fluctuations in pelagic flounder early life stages that have occurred over the past 30 years. We evaluate the hypothesis that the available reproductive volume (habitat), defined as the water column with dissolved oxygen larger than 1 ml/l and salinity between 10.6 and 12 PSU, affects the survival of flounder ichthyoplankton and determines recruitment success.

Both reproductive volume and spawning stock biomass were significant factors determining flounder ichthyoplankton abundance. However, recruitment did not correlate to the supply of larvae what could indicate that important bottlenecks are in nursery grounds in the coastal habitats.

Costal habitats (sandy beaches) are important nursery grounds for pelagic and demersal flounder of Central part of the Baltic Sea. The data for beach seine surveys shows that the smallest flounder juveniles in the end of the summer lives in shallow waters while in the autumn they slowly migrate to deeper waters. The preliminary results indicated two different peaks in the length distribution of flounder juveniles what could be two types (pelagic and demersal) of flounder. Using genetic and otoliths microchemistry analyse it is planned to determine exact proportion of those two types.

Since 2004 an invasive fish species round goby is registered in the coastal waters of Latvia. In the study area (sandy beaches, Pape – Jurmalciems, close to Latvian – Lithuanian border) round goby for the first time was observed in 2009. No changes of flounder recruitment was observed since introduction of round goby while reproduction success of turbot was significantly lower in the last years.

Abstract: Rohtla, M., Vetemaa, M., Svirgsden, R., Taal, I., Saks, L. and Verliin, A.

Spawning habitat preference of Väinameri pike, ide and burbot

Spawning habitat preference (i.e. fresh or brackish water) was investigated in brackish water living populations of pike (Esox lucius), ide (Leuciscus idus) and burbot (Lota lota). Otolith Sr:Ca profiles were used to determine the hatching biome. Of all the adult pike sampled around the Väinameri Sea area (n=435) a total of 90% hatched in fresh water and only 10% in brackish water. These results suggest that brackish water spawning pike may not be as common in the Väinameri Sea as they were suggested to be before the 1970s. Deterioration of brackish water spawning grounds due to the negative effects of eutrophication could be the factor behind low brackish water recruitment. Of all the ide collected from three sites in the Väinameri Sea (n=111) 72% hatched in seasonally freshwater bays (SFB) and only 28% in rivers. The importance of true lotic spawning has decreased substantially in Saunja (most fish hatched in SFB; stocks in the lows) and Käina Bays (all fish hatched in SFB; abundant ide stock). Only Matsalu Bay ide were mostly lotic spawners (stocks in the lows). Of all the burbot collected from the Väinameri Sea (n=74), 96% hatched in fresh water and 4% showed signs of hatching in an environment with Sr:Ca slightly higher than the freshwater threshold, possibly in river mouths. It is concluded that burbot is a fully freshwater spawner in the Väinameri Sea area.
Abstract: Timo Arula

Clupeid spawning ground surveys in the NE Baltic Sea

In this talk results of the four different surveys were discussed: long-term spring herring larval surveys in NE Gulf of Riga (GoR), spring herring spawning ground mapping in Estonian coast, autumn spawning herring in Gulf of Riga and Baltic sprat eggs and larval surveys in the Baltic proper.

NE part of the GoR has been the most important spawning and nursery ground for GoR herring population, where larval herring and related environment was regularly studied since 1947. Larval herring samples were collected weekly from May to August with Hensen larval net and samples were preserved in 4% formalin seawater solution. Individuals were counted and digitalized in lab conditions using zooscanner and later measured from images. We found that the abundance of 1-year old spring spawning herring is statistically significantly $(r^2 = 0.74)$ determined by the number of postflexion herring larvae in the Gulf of Riga (Baltic Sea) in 2004–2013. The abundance of postflexion larvae, in turn, displayed dome-shaped relationship with sea surface temperature experienced since hatching ($r^2 = 0.69$). Winter air temperatures substantially affected seasonality of herring larvae, but not their prey, being thus important factor regulating temporal overlap between them. Most importantly, we have identified a new bottleneck - too high temperature that probably exceeds physiological thermal optimum – for the survival of postflexion herring larvae in shallow estuarine habitats. Future climate warming points to a risk for a decrease in larval herring survival, which may lead to reduction in herring stocks relying in recruitment from shallow coastal areas.

In the 2011–2014 Baltic herring spawning grounds around Estonian coast were mapped using number of different methods, like: underwater bottom recording to identify bottom substrate, hydrology, exposure to the waves, depth e.g. Using recorded data and based on scientific and expert knowledge, MaxEnt (Maximum Entropy Modeling) was applied to create predictive distributional map of herring spawning areas around the Estonian coast.

Baltic autumn spawning herring landings constituted up to 47% in total herring landings in GoR, but dropped drastically to <1% in most recent decades. There has been slight signs on recovery of autumn spawners in GoR. In 2009–2012 historically important spawning and larval herring nursery ground were visited in GoR and distribution and abundance of larvae were compared with historical data. It appeared that the same nursery ground are in use as well as the number of larvae is comparable what it was in 1960s and 1970s before population collapsed. Baltic sprat eggs and larvae distribution was studied in two consecutive years in Baltic proper (Estonian and Latvian EEZ) using Isaac-Kidd larval fish net. Altogether 68 stations were visited in May and June, to collect sprat eggs and larvae towing the net horizontally near the sea surface and vertically from 130m until surface. In each station, CTD was launched to measure water temperature, salinity, oxygen, chl *a*, turbidity from surface to sea bottom. It appeared that bulk of sprat eggs distributed near the south border of Latvian sea zone, and fewer eggs were found in the western part of the Gulf of Finland. In addition, most of the sprat larvae were found from the southern part of Latvian water's and there was no sprat larvae in Estonian waters.

Abstract from *Theme 2* – How are essential habitats for fish mapped and monitored? Methods, available maps, are there data for maps?

Abstract: Meri Kallasvuo

Using high-resolution species distribution modelling to produce reproduction habitat maps of coastal fish to support marine spatial planning

Productive commercial and recreational fisheries are strongly linked to the ecological state of the habitats. Especially the early life stages of fish are very sensitive and the reproduction habitats are usually the most limited habitat type. In the northern Baltic Sea the fish reproduction in most cases takes place in the heavily exploited coastal zone. Therefore in the Finnish national VELMU underwater habitat inventory programme (2004–2015), systematic mapping of the coastal fish reproduction habitats has been conducted. The VELMU programme covers the entire Finnish archipelago with 46,000 km of shoreline. Large spatial and temporal environmental gradients are typical for the extensive archipelago area. In the VELMU programme we have developed and tested new field survey and mapping methods and produced modelled distribution maps of the key reproduction habitats of the most important coastal fish species. These maps serve as basic reference data and allow visual and numerical comparison of coastal areas. They provide concrete support to fisheries and environmental managers. Especially useful these maps have been when planning coastal areas (MSP) and setting local or national fishing restrictions. Also the developed cost-effective field sampling methods

have been useful when consults etc. conduct more detailed surveys for local planning. Main challenges have arisen from the high annual variation and sufficiency of field survey data. The detailed archipelago and large environmental gradients require high-resolution environmental data, of which availability is a challenge. Effort has also been put into building a VELMU online map service to openly share the data and knowledge in as easy-to-understand-form as possible.

Abstracts from *Theme 3* – Conservation and threats on essential coastal habitats for fish

Abstract: Patrik Kraufvelin

Physical threats to essential coastal habitats for fish

In addition to ongoing global environmental change (climate change, eutrophication, overfishing, invasive species, etc.), there is a multitude of physical threats to essential coastal habitats for fish. This presentation focuses on the major physical threats, the kind of disturbances/stresses they impose and their implications for coastal fish habitats. Among the worst physical threats, there is marine ship/both traffic and its infrastructure, coastal engineering measures in bays/on shores and trawling. Less serious or more local threats are imposed by road traffic and its infrastructure, marine extractions, marine renewable energy production, pipelines and cables, vegetation-changing activities and recreation. The different physical threats may disturb fish habitats through direct substrate destruction/loss, erosion, increased sediment cover and turbidity, oxygen deficiency, temperature and salinity changes, noise, changes in water movement/circulation and water level and release of nutrients and pollutants. Most of these stresses are more serious to fish reproduction and juvenile stages, while adult fish sometimes can escape the problem. Noteworthy, the total impact of small boats may be worse than ship traffic due to the high frequency of the former (it is occurring everywhere, also in shallow waters) and because small boat traffic is most intense during the biologically most active and sensitive summer period. Socio-ecologically it is evident that also man will be affected if key-habitats for fish disappear and many ecosystem services are indeed threatened by anthropogenic physical activities.

Abstract: Ulf Bergström

Essential fish habitats in management

Management of fisheries and nature conservation has historically been separated in the Baltic Sea region, and the awareness of potential synergistic effects between the two has been low. Recently, several marine ecological studies have shown how mutual benefits may be reached through an integrated management of fisheries and habitats. Healthy vegetated habitats are needed for the reproduction of many coastal fishes, while strong populations of predatory fish may through maintaining trophic cascades support habitat-forming vegetation. To increase the awareness of the benefits of integrating management of fisheries and habitat, the scientific community may contribute by: 1) demonstrating the ecological synergies that may be achieved by protecting essential fish habitats (EFH), 2) developing methods for and carrying out large-scale mapping of EFH, 3) quantifying the effects of different threats to these habitats, and 4) communicating the importance of the habitats.

Habitat maps are produced by extensive surveys, usually combined with statistical modelling, or, as a first and simpler approach, by interview studies. The maps may be used in marine spatial planning to protect sensitive areas from threat like shoreline development, dredging, boating etc. They can also be used to evaluate and strengthen MPA networks. To quantify threats to EFH, maps of the pressure variables, together with a mechanistic understanding of the effects of different threats on the habitats, are needed in addition to habitat maps. Having this information at hand, it is possible to perform scenario analyses to explore the effects of different management regimes. Similar analyses may be useful in marine spatial planning for efficient use of space. The importance of EFH may be communicated through valuation of ecosystem services, such as food, recreation, biological regulation. An efficient way of increasing the awareness of these sensitive coastal environments and the will to protect them is to reveal their beauty visually through photos and videos.

Abstract: Antti Lappalainen, Meri Kallasvuo

Conservation and threats on essential coastal habitats for fish – Finland

In Finland, three typical reproduction habitats for coastal fish exist: shallow sandy beaches, flads and inner bays and estuaries are. 1) Shallow sandy beaches are important habitats for sea-spawning whitefish larvae and YOY flounder and turbot. Sandy beaches are protected by Nature Conservation Act. Eutrophication forms the main threat causing accumulation of drifting filamentous algae and in extreme cases, even slow transformation into reed belt shores in sheltered areas. 2) There are around 10,000 flads or small lagoons in Finland. They are important reproduction habitats for perch and pike, especially in the Gulf of Bothnia. Small pristine flads (<10 ha) are protected by Water Act and flads in Natura 2000 areas protected by Habitat Directive. Land lifting is a natural threat to flads especially in the Quarken area, but also various coastal activities and eutrophication endanger these habitats. 3) Inner bays and estuaries are main reproduction areas for pike, pikeperch, cyprinids, perch and burbot. These habitats are generally not very strictly protected. Inner bays and estuaries are a common habitat type, especially in the southern and southwestern Finland. As a whole, these habitats are perhaps not yet seriously threatened. However, eutrophication and climate change are altering also inner bays and estuaries.

To conclude, Finnish coast consists of diverse coastal reproduction habitats. Eutrophication and climate change threaten quality of the habitats. Anthropogenic activities, such as dredging and coastal constructions, are more or less local problems. Conservation by protected areas is implemented on several levels: e.g. by Natura 2000, private conservation areas and fishing restrictions during spawning time.

Abstract: Elliot John Brown

Conservation and threats on essential coastal habitats for fish: The need for protection of essential fish habitats

The tension between the protection of productive, diverse near-coast, marine habitats and the pressures placed upon them by largely coastal human populations is well documented. This conflict is often referred to by natural scientists when introducing specific investigations into anthropogenic impacts. The conclusions from these studies often use the effects on fish populations and fisheries catches as evidence of the need for changes to regulatory protection. However, to effectively communicate this, one must consider the value that society at large places on both fish stocks and the resources that their habitats provide. This presentation investigates the value of fisheries and the supporting habitats to humans in a broader context than fisheries data and uses specific examples with the intention to initiate discussion on the need to protect essential fish habitats from a diversity pressures. Examples of values include: (1) The effects of commercial fishery collapses and closures on regional economies and the individuals within. (2) Non-commercial catches – how they can be undervalued and their scale. Physical habitat as a resource; (3) rock, sediment, minerals (4) seawater and (5) space. (6) Finally, the indirect value given to the knowledge of an improved and healthy environment. In summary, as natural scientists we should consider all of the ways in which coastal fish habitats provide value to society and use these as examples when communicating the need for the protection of essential fish habitats.

Abstract: Linas Ložys and Justas Dainys

Essential coastal habitats for fish in Lithuania: Threats and conservation

The Baltic Sea coastline of Lithuania is only about 100 km long. Freshwater Curonian Lagoon is connected with the Baltic Sea through Klaipeda port channel. It is the main gate for fish migrations between the Baltic Sea, Curonian Lagoon and further upstream to the basin of the Nemunas River or *vice versa*. Few abundant fish species are spawning in Lithuania's coastal waters: herring (*Clupea harengus*), flounder (*Platichthys flesus*), turbot (*Psetta maxima*) and round goby (*Neogobius melanostomus*) – recent invader of Ponto-Caspian origin. Perch (*Perca fluviatilis*), ruff (*Gymnocephalus cernuus*) and most *Cyprinid* species spawn in the Lagoon, while pikeperch (*Sander lucioperca*), smelt (*Osmerus eperlanus*), burbot (*Lota lota*),

vimba (*Vimba vimba*), salmon (*Salmo salar*) and sea trout (*Salmo trutta*) migrate for spawning to Nemunas River and tributaries.

The bottom structure of Lithuania's coastal waters is mapped roughly long decades ago. More detailed maps are developed during the recent decade as the result of various research projects. E.g. multibeam bathymetry maps and seabed profiles were developed as part of PhD project at Klaipeda University aiming to detected herring spawning beds; the study revealed most essential herring spawning sites in the areas of bottom elevations and their slopes in the Lithuania coastal waters. As the result of recent LIFE-NATURE project (DENOFLIT) twaite shad and whitefish (Coregonus lavaretus) abundance in trawling surveys in the Baltic Sea in Lithuania's EEZ during 1995–2010 was mapped and it suggests most essential habitats for these species at the Baltic Sea. Recent decline of twaite shad and whitefish stocks is presumably related to overfishing at spawning sites in the Curonian Lagoon. Such threats as Cormorant predation on whitefish juveniles, eutrophication and change in temperature regime at spawning sites may also be related to the decline of whitefish population. Mapping of *Salmonids* migration through the Curonian Lagoon is based on the detail analysis of fishery logbooks' daily data. The mapping done allows improving protection of Salmonids by closing fishery in most essential for the species migration areas during particular periods. Cormorants are often blamed as one of most essential threats for fish stocks in the Curonian Lagoon and coastal waters. However, monitoring of fish stock and stabile landings in the fishery suggest minor effect of Cormorants on the local fish populations. At the same time Cormorants play positive role in ecosystem recently by controlling of round goby abundance in the coastal waters and the Curonian Lagoon. Abundance of round goby especially peaked in 2011 after first record of the species in Lithuania in 2002. This invasive species is evidently among most essential threats for the ecosystem in coastal waters. Round gobies dramatically reduced numbers molluscs, such as Mytilus edulis. Mytilus edulis are habitat-forming species on soft and especially on hard bottoms, their recent decline as the result of Round goby predation results in loss of overall bottom habitat diversity. Decline of *Mytilus edulis* and associated species makes essential threat for many other fish species as well as wintering birds. A decline of some wintering bird species in the area is hypothetically associated with effects of decline in invertebrate species typically used by birds during wintering period.

Abstract: Martin Snickars

Coastal lagoons - vulnerable nursery habitats

In many coastal waters of the Baltic Sea, shallow vegetated areas constitute essential fish habitats for fish species, such as perch, pike and cyprinids, which use these habitats for spawning, nursery and feeding. Coastal lagoons, e.g. silled flads, are one type of nursery habitats. Naturally, silled lagoons become gradually isolated from the sea due to the on-going landuplift. The isolation process creates beneficial nursery habitat as temperature and vegetation cover increase with topographic isolation, and continues to do so as long as the sill area is unaltered.

However, shallow, wind-protected bays and lagoons are also attractive sites for marinas and recreational activities, which may compromise the function of these habitats. Major threats are dredging, boating, and infrastructure. Studies have shown that anthropogenic stressors are equally important as environmental variables, and that coastal lagoons in natural conditions are becoming rare. The results highlights that the species composition of vegetation sensitive vs. tolerant species) affects the abundance of juvenile fish. The abundance increases with the cover of sensitive species, suggesting that pressures affect the ecosystem functioning by altering the vegetation community. Effective conservation of nursery areas is important as many shallow vegetated habitats are vulnerable to major threats of coastal areas.

Abstracts: Peter Möller

Bottom habitat measures within the coastal areas of the State of Mecklenburg-Western Pomerania

It is not allowed to use bottom trawls within the 3 nm zone off the Mecklenburg-Western Pomerania coast and in the inner parts (lagoons). Exceptions are catch of live bait for personal use by long-line fishing (rather seldom) and for touristic activities, the use of trawls with drifting (sailing) boats (Zeesenfischerei, zeese fishery by one or two boats) in the inner parts of the coast (e.g. in the Darß-Zingst Bodden Chain). Fishermen have to apply for permission.

There are different government acts (State of Mecklenburg-Western Pomerania and EU Commission) to protect several marine and terrestrial areas (e.g. Bird Directive, Habitat Directive, Water Framework Directive, Maritime Spatial Planning of the State M-WP (see map or Regierungsportal www.regierung-mv.de). Within National Parks (e.g. Nationalpark Nordvorpommersche Boddenlandschaft), no bottom disturbing activities (e.g. drilling or sand extraction) are allowed.

In the Baltic Sea Exclusive Economic Zone (EEZ), 5 Natura 2000 areas (Flora Fauna Habitat Directive (92/43/EWG): Adlergrund (DE 1251-301), Fehmarn Belt (DE 1322-301), Kadetrinne (DE 1339-301), Pommersche Bucht mit Oderbank (DE 1652-301), Westliche Rönnebank (DE 1249-301) and 1 Special Protected Area (Bird Directive (79/409/EWG): Pommersche Bucht (DE 1552-401) exist. There is still an ongoing discussion between the Federal Ministry of Environment and the Ministry of Agriculture about a management plan in these areas concerning fishing activities. A sediment map can be found at www.geoseaportal.de. There are several seasonal closed spawning areas within the inner parts of the coastal area where fishing is forbidden at main spawning season, but this is not a "habitat" protection measure. Every activity in the coastal zone is regulated by different acts.

Projects of the Institute of Fisheries, State Research Centre of Agriculture and Fisheries and the Research Association Fish and Environment

Eel project: The EU required data from all member states about the situation of the eel stock to develop a new management plan for eel (2007). To collect data about the density of eel in the coastal waters of Mecklenburg-Western Pomerania an enclosure sampling gear was developed by our fisheries engineer (Research Association Fish and Environment). The transportable fishing system consists of two parts. The external part consists of a boundary net arranged in a 100 m × 100 m square so as to enclose an area of 1 ha. The mesh size of the lead line weighted boundary net is 10 mm and net height is 1.8 m. The boundary net has fyke net chambers (mesh size 11 mm) at each at its four corners. Additionally, six chains of fyke nets (in total 4 double chamber fyke nets for one chain, 3 funnels in each chamber and 40 cm high leaders) with a mesh size of 10 mm in the end chamber are deployed to increase the likelihood of catching as many eels as possible within the boundary net. In total, 200 samples were taken in 2008–2011 (see Ubl and Dorow et al. 2014). From 2011 on, some of the sampling areas were completely recorded with sonar. It was a test to correlate the number of caught eel with the structure of the bottom.

Artificial Reefs: The reef area is more or less 1 nautical mile out on the sea, in front of the summer resort area of Nienhagen (Mecklenburg-Western Pomerania). First, a project study was started from 1994–1996 and small structures were installed in 1996. 2,000 tons of natural stones were added in 1998. The major project started in 2002. In 2003, concrete elements were added to it and the reef area grew up to 4 hectare and finally

a research platform, for measurement and a radio connection to send data to a terrestrial station, was built in 2010. Several projects took place in this area. Currently, the final reports are being prepared. It is a kind of man made refuge for fish. And that causes very heated debates about the pros and cons of artificial reefs. Some people think it is a useful tool as a measure to outbalance construction projects and create refuges for several species. Others see that as a far too strong influence or change in the bottom habitat.

Abstract: Saks, L., Vetemaa, M., Taal, I., Verliin, A. & Rohtla, M.

Coastal marine habitats and their conservation status in Estonia

Coastal habitats of Estonia are mapped by modelling exercises in which direct "point data" and indicator species serve as model inputs. This has been done for soft sediments, sandy bottoms, sandbanks, reefs, "tidal" mudflats and sandflats. Large part of the Estonian coastal sea is relatively well protected, mostly as Natura 2000 nature protection areas. All these areas have different restrictions which apply in various circumstances. For example, different fishing restriction can be now easily found at: http://pump.regio.ee/kalandus/public/

Abstract: Zusana Celmer

Coastal fish habitats - the Puck bay case

The Puck Bay is one of the most important coastal habitats for fish in Polish waters. The water body is very shallow and therefore amplitudes of water temperature are high. Water exchange is difficult due to occurrence of underwater sandy reef. The basin provides suitable conditions for feeding and spawning for many fish species.

In the 60s the Puck Bay was in a healthy condition. Probably, due to direct discharge of sewage from the surrounding area, the state has deteriorated markedly. The majority of vascular plants were replaced by filamentous algae (bladderwrack disappeared). Feeding, reproduction and also nursery grounds for larvae and adults of some species were lost. In recent years, the situation has markedly improved as the consequence of the activity of sewage treatment plant. Although seagrass meadows slowly reborn, it seems unlikely to achieve the environment observed 60 years ago. Since 2004, the Puck Bay has been protected under the Natura 2000 (Habitats and Birds directive). The current major threats are unregulated tourism, fishing poaching, excessive growth of filamentous algae, high abundance of stickleback and unbalanced development of the maritime infrastructure.

Appendix 4 – Information from Baltic Sea countries about mapping and monitoring of essential fish habitats

Table 6: Denmark

Spec mapj moni	ies that are ped and itored	Observation / Monitoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables collected	Geographical cov- erage and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
Cod, brill, sole	plaice, turbot, eel, lumpfish,	Combination of fisher interviews, VMS data for ves- sels > 12 m, IBTS data	GIS layers-map, overlay-maps on sediment, depth maps combined with fish presence maps	For fisher inter- views: Murray <i>et al.</i> 2008ab; Neis <i>et al.</i> 1999; Gunnartz <i>et al.</i> 2011	Habitat variables used: sediment maps, depth maps, macroalgal coverage	The Sound 2015	Provide fish habitat maps for manage- ment of marine ag- gregate extraction	GIS maps, interview notes, VMS data, Fish Atlas data (available after publication)	DTU Aqua (fish VMS and fisher in- terview), Zoological museum (Fish Atlas), GEUS (sediment maps); Thomas Kirk Sørensen
20 sp moni	pecies itored	International stand- ardised fish moni- toring	From monitoring data and other in- formation, maps of spawning areas, ju- venile areas and nursery areas from different time peri- ods of different species	Warnar <i>et al</i> . 2012	No information	All Danish waters	Baseline for MSFD	Warnar <i>et al</i> . 2012.	DTU Aqua; Josianne Støttrup
All sp in gil fyker	pecies caught Inets or nets	Standardised fykenets or one mesh-size gillnets. Fishermen registra- tions	No mapping as yet	HELCOM GUIDE- LINES (modified) and latest report: Kristensen <i>et al.</i> 2014	Temperature loggers	All inner Danish wa- ters	Year round sam- pling (fish sampled monthly, year- round unless ice cover, data regis- tered by recrea- tional fishermen)	No maps yet	DTU Aqua; Josianne Støttrup

Table 7: Estonia

Species that are mapped and monitored	Observation / Mon- itoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables collected	Geographical coverage and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
23–36 species	Standardised coastal fish moni- toring routines for coastal areas of the northern Baltic Sea, with some modifications	No mapping as yet	HELCOM GUIDELINES (modified)	Secchi depth, temperature	Matsalu Bay 1993–2014; Vaindlo 1997–2014; Saarnaki 1998–2014; Vilsandi 1993–2014; Käsmu Bay (Eru Bay) 1997–2014; Kihnu 1997–2014; Kõiguste 2005–2014; Küdema Bay 2000–2014; Pärnu Bay 2000–2014	Coastal fish monitoring	Excel files	EMI; Markus Vetemaa
10–29 species	Standardised coastal fish moni- toring routines for coastal areas of the northern Baltic Sea, with some modifications	No mapping as yet	HELCOM GUIDELINES (modified)	Secchi depth, temperature	Väike väin 2008; Vilsandi 2009; Aksi 2009; Häädemeeste 1999 and 2002; Heltermaa 1998, 2009– 2010; Käina Bay 2005, 2010; Kõpu 2011; Lahepera Bay 2013; Laidevahe 2000; Naissaar 2004– 2005, 2009–2010; Neugrund 2007–2008; Narva Bay 2007– 2014; Pakri 2009; Paldiski 2001– 2002; Prangli 2004–2006, 2009; Rohuküla 2009–2010, 2014; Ruhnu 2008; Kuressaare Bay 1999, 2002; Gulf of Finland (Tal- linna madal etc.) 2012; Suur väin (Great belt) 2008–2009; Tamme (Küdema Bay) 2005–2007; Virtsu 1993, 1999, 2009–2010; Uudepanga Bay 1999; Vormsi 2000; Matsalu bay 2001–2002. Käsmu bay 2003–2004	Inventory (Year- round sampling was undertaken in Matsalu bay during 2001– 2002 and in Käsmu bay dur- ing 2003–2004. See e.g. Vetemaa <i>et al.</i> 2006 for details)	Excel files	EMI; Markus Vetemaa
13 species	Standardised coastal fish moni- toring routines for coastal areas of the northern Baltic Sea were followed	No mapping as yet	HELCOM GUIDELINES	Water chemical properties, salin- ity, macrophytes, zooplankton (1993)	Saunja 1993?, 2005	Inventory	Excel files	EMI; Markus Vetemaa

Table 8: Estonia (continued)

Species that are mapped and monitored	Observation / Mon- itoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables collected	Geographical coverage and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
20 species	Multimesh gillnets (nordic coastal)	No mapping as yet	BONUS IN- SPIRE, man- ual	Secchi depth, temperature, sa- linity, bottom type, oxygen	Dirhami 2014	Inventory	Excel files	EMI; Markus Vetemaa
8 species	Lacustrine gillnets	No mapping as yet	HELCOM GUIDELINES (modified)	Secchi depth, temperature	Sutlepa Sea 2005	Inventory	Excel files	EMI; Markus Vetemaa
13–26 species	Beach seine	No mapping as yet	Taal <i>et al.</i> 2014	Temperature	Eru Bay 2008–2009; Mõntu 2004–2006; Saarnaki 2004–2005; Ruhnu 2004; Parasmetsa Bay 2004–2005; Panga 2004–2006; Pakri 2006; Kõiguste 2006	Inventory (Fish were sampled monthly during ice-free period from April 2008 to December 2009 only in Eru bay)	Excel files	EMI; Markus Vetemaa
Baltic herring (Clupea harengus)	Field work com- bined with remote sensing data	MaxEnt (Maximum Entropy Modeling), raster dataset on environmental vari- ables (based on sci- entific and expert knowledge)	No reference	CTD data	All Estonian coast 2011–2014	Map Baltic her- ring spawning grounds in Esto- nian coast	GIS predictive dis- tributional maps	EMI; Jonne Kotta
Baltic herring (Clupea harengus)	Field work com- bined with remote sensing data	Modeling, raster dataset on environmental vari- ables	Saat 2015	CTD data	All Estonian coast 2011–2014	Map Baltic her- ring spawning grounds in Esto- nian coast	GIS predictive dis- tributional maps	EMI; Jonne Kotta

Table 9: Estonia (continued)

Species that are mapped and moni- tored	Observation / Monitoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables collected	Geographical coverage and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
Benthic macro- phyte communities	Field work com- bined with remote sensing data	MaxEnt, raster da- taset on environmental variables	HELCOM 1998	Algae, aquatic plants, bottom animals by scuba- diving, drop-vid- eos, grab samples etc.	All Estonian coast 2005–2013	Habitat mapping	GIS predictive dis- tributional maps	EMI; Georg Martin
Distribution of habitats and key species In Estonian territorial waters	Modelled by using direct "point data" and by using indi- cator species	Modeling, raster dataset on environ- mental variables	Herkül 2014	Algae, aquatic plants, bottom animals, etc.	All Estonian coast 2005–2013	Habitat mapping	GIS predictive dis- tributional maps	EMI; Kristjan Herkül

Table 10: Finland

Species that are mapped and monitored	Observation / Mon- itoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables collected	Geographical coverage and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
Newly hatched lar- vae of pike, roach (other cyprinids), (burbot), (stickle- backs)	White plate	Species distribution modelling by lo- gistic regressions	Kallasvuo 2010	VELMU program: Algae, aquatic plants, bottom animals by scuba-diving, drop videos, grab sam- ples, environmental background data	Vegetated shores along the Finnish southern and southwestern coast 2004–2010 (May–July)	Habitat map- ping for man- agement and MSP	Freely available 01/2016 in a map portal	Luke; Meri Kallasvuo
Newly-hatched lar- vae of whitefish, vendace	Beach seine	Species distribution modelling by Gaussian processes	Vanhatalo <i>et</i> <i>al</i> . 2012, Ve- neranta <i>et</i> <i>al</i> . 2013	VELMU program and In- tersik / Norrsik: Algae, aquatic plants, bottom animals by scuba-diving, drop videos, grab sam- ples, environmental background data	Shallow sandy beaches along the Finnish coast 2009–2010, 2013 (April–May)	Habitat map- ping for man- agement and MSP	Freely available 01/2016 in a map portal	Luke; Meri Kallas- vuo and Lari Vene- ranta
YOY/juveniles of flounder, turbot	Beach seine	Only point data, no modelling	Wijkmark et al. 2014	VELMU program: Algae, aquatic plants, bottom animals by scuba-diving, drop videos, grab sam- ples, environmental background data	Shallow sandy beaches along the Finnish south- western coast 2010– 2015 (May, September– October)	Habitat map- ping for man- agement and MSP	Freely available 01/2016 in a map portal	Luke; Meri Kallas- vuo and University of Helsinki; Henri Jokinen
European flounder, (turbot)	Beach seine	Point data	Jokinen <i>et</i> al. 2016	No	Four (five) sites on the Åland Islands and two sites at the Hanko penin- sula in two periods: 1979–1992 and 2012– 2014 (biannually: late October and early May)	To assess tem- poral changes in juvenile floun- der abundances	Available upon re- quest	Luke (1979–1992 data), H. Jokinen (Eero Aho) / Tvärminne zoologi- cal station, Univer- sity of Helsinki (2012–2014 data)

Table 11: Finland (continued)

Species that are mapped and monitored	Observation / Mon- itoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables collected	Geographical cover- age and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
European floun- der and turbot (+ Pomatoschis- tus sp., Ammo- dytes, Gas- terosteidae, other littoral species)	Beach seine	Point data	Epifauna with drop trap sam- pling (Nohrén <i>et</i> <i>al.</i> 2009);	Sediment grain size, or- ganic content, salinity, pH, turbidity, vegetation / algae cover and bio- mass, infauna, depth, ex- posure index	21 sites around Hanko peninsula in mid-August 2013	Investigation of cur- rent occurrence of ju- venile flatfish, charac- terization of juvenile flatfish habitats and as- sessment of juvenile occurrence in relation to vegetation / algae and other habitat char- acteristics	Available upon request	Tvärminne zoologi- cal station (Univer- sity of Helsinki); Henri Jokinen
Newly-hatched larvae of Baltic herring, pike- perch, perch, smelt, (gobies)	Gulf ichthyoplank- ton samplers	Species distribution modelling by Gaussian processes	Veneranta <i>et al.</i> 2011, Kallasvuo <i>et al.</i> in prep.	VELMU program: Algae, aquatic plants, bottom animals by scuba-diving, drop videos, grab sam- ples, environmental background data	Entire Finnish coastal area 2007– 2014 (May–June)	Habitat mapping for management and MSP	Freely available 01/2016 in a map portal	Luke; Meri Kallasvuo
YOY/juveniles of pike, perch, cy- prinids, sticklebacks, species with swimbladder	Detonations	Mainly point data, but species distri- bution modelling (GAM, MARS, RF) has been con- ducted using deto- nations in the Ar- chipelago Sea	Snickars <i>et al.</i> 2007	VELMU program: Algae, aquatic plants, bottom animals by scuba-diving, drop videos, grab sam- ples, environmental background data	Vegetated shallow bays in SW Finland and Åland archipel- ago 2002–2004, 2006 (July–Septem- ber)	Habitat mapping for management and MSP	Available by re- quest	Luke; Meri Kallas- vuo and Åbo Aka- demi University; Martin Snickars
Trials on other larvae/juvenile fish	Dipnet, pushnet, scuba-diving, gillnets	Only point data, no modelling	Not available	VELMU program: Algae, aquatic plants, bottom animals by scuba-diving, drop videos, grab sam- ples, environmental background data	Entire Finnish coastal area 2004– 2014 (April–Septem- ber)	Habitat mapping for management and MSP	Available by re- quest	Luke; Meri Kallasvuo

Table 12: Germany

Species that are mapped and monitored	Observation / Mon- itoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables col- lected	Geographical cover- age and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
European eel, non-target spe- cies	Underwater video cameras, sonars, eel sampling system	Surfer Golden Soft- ware	Non, newly de- veloped	Water temperature, salinity, pH, oxygen level, visibility, wind, weather, currents, scuba diving photo and video	Coastal areas M-WP, since 2010	Correlation habitat and abundance	Yes, XYZ data shapes	Federal research centre M-WP and Fish and Environment; www.verein@fischumwelt.de, Peter Möller, Olaf Krüger
Whitefish, non- target species	Underwater video cameras, trawl net (zeese), beach seine	Surfer Golden Soft- ware	HELCOM GUIDELINES (modified)	Water temperature, salinity, pH, oxygen level, visibility, wind, weather, current	Coastal areas and lakes M-WP, Since 1995	Stock assess- ment, spawning areas, repro- duction	Yes, XYZ data shapes	Fish and Environment; www.verein@fischumwelt.de, Thomas Lorenz
Cod, non-target species on artifi- cial reefs	Underwater video cameras, tags (inside & outside), gillnets, fish traps	Surfer Golden Soft- ware	HELCOM GUIDELINES (modified)	Water temperature, salinity, pH, oxygen level, visibility, wind, weather, current, scuba diving photo and video	Coastal areas M-WP, since 2002	Local stock as- sessment, food web, migration	Yes, XYZ data shapes	Federal research centre M-WP and Fish and Environment; www.ver- ein@fischumwelt.de, Bodo Dolk, Olaf Krüger, Peter Möller
Seatrout	Underwater video cameras, tags (inside & outside), gillnets, electro fish- ing	Surfer Golden Soft- ware	HELCOM GUIDELINES (modified)	Water temperature, salinity, pH, oxygen level, visibility, wind, weather, current, scuba diving photo and video	Coastal area and creeks M-WP, since 1993	Stock assess- ment, spawning areas, repro- duction	Yes, XYZ data shapes	Federal research centre M-WP and Fish and Environment; www.verein@fischumwelt.de, Harry Hantke
Fishes of the Ad- lergrund	Underwater video	Videos	HELCOM GUIDELINES (modified)	Water temperature, salinity, pH, oxygen level, visibility, wind, weather, current, scuba diving photo and video	EEZ Germany	Fish composi- tion of the Ad- lergrund	Yes, XYZ data shapes	Palaemon Aquatic service; www.verein@fischumwelt.de, Thomas Lorenz

Table 13: Germany (continued)

Species that are mapped and monitored	Observation / Monitoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables col- lected	Geographical cover- age and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
Fishes of the Szczecin Lagoon and Peenestrom	Underwater video, egg sampling	Maps	HELCOM GUIDELINES (modified)	Water temperature, salinity, pH, oxygen level, visibility, wind, weather, current, scuba diving photo and video	Szczecin Lagoon and Peenestrom, 1997– 1998	Spawning grounds	Yes, XYZ data shapes	Fish and Environment; www.verein@fischumwelt.de, Thomas Lorenz
Flounder, perch, pikeperch, round goby etc.	Bottom trawling; eel trawl since 2012 (bottom ot- ter trawl OTB- TV3-520)	Abundance and bi- omass per area at different sampling sites	No reference	Temperature, oxygen, salinity, bot- tom type	Pomeranian Bay be- tween Usedom island and Oderbank 2003– 2015	Monitoring of fish biodiversity	Access data- base	German Sea fishery institute, University of Rostock; Winkler Helmut and Böttcher Uwe
YOY fish (smelt, gobies,etc.)	Special bottom trawl for small fish	Abundance and bi- omass per area at different sampling sites	No reference	Temperature, Oxy- gen, salinity, bottom type, nutrients, zoo- plankton	Darss-Zingst lagoon 1998–2015	Monitoring of YOY fish bio- mass	Exel database	University of Rostock; Winkler Helmut

Table 14: Latvia

Species that are mapped and monitored	Observation / Monitoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables collected	Geographical cov- erage and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
Round goby	Gillnets, beach seine, video, scuba	Not available	No reference	Vegetation, Bottom type	Latvian coast 2015–2016	Round goby distri- bution in Latvian coast	Not yet	BIOR and LHEI; Di- dzis Ustups and Strake
Flounder	Beach seine	Not available	BONUS INSPIRE, manual	Vegetation, Bottom type	Irbe Strait 2014– 2015	Distribution of flounder juveniles	Not yet	INSPIRE and BIOR; Didzis Ustups
Flounder, cod	Gillnets	Not available	BONUS INSPIRE, manual	Vegetation, Bottom type	Latvian coast, SD 28, 5 to 50 m tran- sect (one location) 2014–2015	Spatial planning	Not yet	INSPIRE and BIOR; Didzis Ustups
Sprat, herring	Hydroacoustic sur- vey	Surfer Golden Soft- ware	No reference	No	Baltic Sea 2004– 2013, Gulf of Riga 2004–2013	Spatial planning	jpg (shape)	BIOR; Didzis Ustups
Cod, flounder	BITS	Not available	No reference	No	Baltic Sea 2004– 2013	Spatial planning	jpg	BIOR; Didzis Ustups
Round goby	Commercial catches, gill net surveys, beach seine	Information from commercial log books are used as input data	Knospina and Put- nis 2014	No	Latvian coastal area 2004–2015	Spatial planning	jpg (shape)	BIOR; Elina Knospina

Table 15: Lithuania

Species that are mapped and mon- itored	Observation / Mon- itoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables collected	Geographical cov- erage and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
Herring	Multibeam bathym- etry, Scuba diving, side-scan sonar, benthic roe samples	Spawning sites marked on multibeam bathymatry map, seabed profiles ana- lysed.	Šaškov et al. 2014	Bathymetry, sedi- ments, Broad BPI (Bentic position in- dex), Slope	Lithuanian coastal waters 2009–2010	PhD-thesis	Maps	Saskov, A. and Linas Lozys
Whitefish, twaite shad, cod, floun- der, others	Bottom trawling	CPUE visualised using GIS	Maps are available at http://corpi.ku.lt/ denoflit/	No	Lithuanian coastal waters and EEZ waters 1995–2010	Inventory of fish distribu- tion in Lithua- nian marine waters	Maps	Lithuanian fishery ser- vice and nature re- search centre; Linas Lozys
Commercially ex- ploited fish spe- cies	Daily and/or monthly records of logbooks	CPUE visualised using GIS	Various project reports e.g. "LITCOAST"	No	Lithuanian coastal waters and the Cu- ronian lagoon 1998–2015	Fishery con- trol and spe- cies distribu- tion research	Maps, excel files	Ministry of environ- ment, Ministry of agri- culture, Nature re- search centre; Linas Lozys
Biological and ge- ological features of bottom habitats	Remote video cam- eras; standard grab sampling; sampling by SCUBA, side scan surveys	Data visualised using GIS	Martin <i>et al.</i> 2010	Bentic fauna, depth, vegetation	Lithuanian coastal waters 2006–2007	Life project	Maps	Daunys, D. and Linas Lozys
Fish juveniles	Beach seine	Data not mapped yet	No reference	No	Lithuanian coastal waters 2005–2007	Life project	Excel files	Linas Lozys
Fish juveniles	Beach seine	Data not mapped yet	No reference	Sediments, macro- phyte abundance, depth, tempera- ture, salinity	Curonian lagoon (Lithuanian part) 2009–2015	Monitoring	Excel files	Linas Lozys

Table 16: Poland

Species that are mapped and monitored	Observation / Mon- itoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables col- lected	Geographical cover- age and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
All	Review and synthe- sis of available data and maps	Different mapping methods	No reference	Sediments, macro- phytes, zoobenthos	All Polish EEZ, with special emphasis to Puck Bay, Slupsk Bank, and Rowy-Ustka coastal area. Archive data, the oldest from 1970s and 1980s.	To start marine habitat mapping in Poland	www.pom-habi- taty.eu/en	?
Only habitat map- ping	Direct observations (macrophytes), sedi- ment samples, mod- ified Bongo net hauls, ortophoto- maps analyses	Basic bathymetry of the shallow coastal zone – point measured depth, analyses of distribution of hard bottom vegetation from orthophotomaps (aerial scanning material). Synthesis based on GIS methods	No reference	Depth, basic sediment analyses (mean grain size, silty-clay con- tent, organic matter), macrophytes (taxon- omy, species compo- sition, distribution)	Szczecin Lagoon 2012– 2015	Research project (Habitat mapping of shallow (coastal) areas of Polish part of the Szczecin Lagoon)	Not available before the end of the project	NMFRI; Adam Woźniczka
Pikeperch, bream	Fish larvae sampled by neuston net (2x1 m, 500 µm). Fish sampled by Nordic survey nets, "regu- lar" gill and fyke nets, beach seine and electrofishing	Mapping of spawning grounds	Official methods accepted ac- cording to WFD monitoring in Poland	?	Vistula lagoon 2010– 2012	Research project (Habitat mapping of shallow (coastal) areas of Polish part of the Szczecin Lagoon)	Limited availa- bility, only via contact person	NMFRI; Dariusz Fey
Herring (larvae)	Bongo net and neus- ton net hauls	Distribution maps based on sampling data (Bongo and neuston net hauls). GIS methods.	No reference	Salinity, temperature, depth	Vistula Lagoon, Pomeranian Bay, since 2007	Research project (Regular monitor- ing of herring lar- vae on important spawn- ing areas)	Not available before the end of the project	NMFRI; Dariusz Fey

Table 17: Sweden

Species that are mapped and monitored	Observation / Monitoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables collected	Geographical cov- erage and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
Turbot, flounder, whitefish, pike- perch, herring, ven- dace	Interview study with commercial fishermen and managers	Delineation of spawning areas on maps	Gunnartz <i>et al.</i> 2011	Species, year, time of spawning, origin of information, spawning indica- tions, substrate, depth	Baltic Proper, Both- nian Sea, Bothnian Bay. Interviews performed 2003	Obtaining full cover maps of spawning areas based on lo- cal knowledge	Freely available (without attributes) from Swedish na- tional authorities	SWaM; Ulrika Gunnartz
Perch, pike, pike- perch, cyprinids, sticklebacks	Underwater detonations	Species distribution modelling	Snickars <i>et al</i> . 2007, Sundblad <i>et al</i> . 2014, Bergström <i>et al</i> . in prep.	Depth, substrate, macrophytes, tem- perature, salinity, turbidity	Baltic Proper, Both- nian Sea, Bothnian Bay. Approximately 11,000 detonations between 2001 and 2014	Juvenile fish sur- veys. Mapping within research projects as well as regional mapping efforts	Variable. Nursery habitats have been predicted in five counties, and are available	Regional authorities; Göran Sundblad, AquaBiota Water Re- search
Whitefish, ven- dace, stickleback	Beach seine	Species distribution modelling	Florin <i>et al.</i> in prep	Substrate, vegeta- tion cover, algal mats, depth, tem- perature, salinity	Baltic Proper, Both- nian Sea, Bothnian Bay	Mapping project	Will eventually be freely available pointdata in SLU database KUL	SLU, HaV, Regional County Boards; Ann-Britt Florin
Flounder, turbot	Beach seine	Only point data, no modelling	Survey Protocol Beachseine surveys BONUS INSPIRE, Nissling <i>et al.</i> 2007, Martinsson 2011	Substrate, turbid- ity, cover of vege- tation, depth, tem- perature, salinity, wave height	Baltic Proper	Research project	2014 and 2015 data freely availa- ble after BONUS INSPIRE project fin- ishes in 2018; older data available upon request to UU	BONUS INSPIRE 2014–2015 and older data from Uppsala University; Anders Nissling, UU

Tab	le 18:	Swede	n (continued)
-----	--------	-------	--------------	---

Species that are mapped and monitored	Observation / Monitoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables collected	Geographical cov- erage and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
Perch	Visual observation of egg strands by snorkelling	Habitat modelling on some of the data	For sampling: Snickars <i>et al.</i> 2010; For produc- tion of maps: Sund- blad <i>et al.</i> 2011; Additional data from: Sundblad <i>et</i> <i>al.</i> 2014	Vegetation (species and cover), depth, temperature, salin- ity, turbidity	Baltic Proper, Both- nian Sea 2003 and 2007	Research project	Available in Swe- den (Regional county board of Stockholm)	Maps from Re- gional county board of Stock- holm; Göran Sundblad, AquaBi- ota Water Research
Herring, sprat, sticklebacks	Hydroacoustics	Species distribution modelling	Nyström Sandman <i>et al.</i> 2013ab	Not available	Baltic Proper 2011	Mapping project	Free at low resolution	Regional county boards in Stock- holm and Söder- man-land; Tomas Didrikas, AquaBiota Water Research
Flounder, cod, tur- bot, round goby	Gillnets	Species distribution modelling	Survey Protocol Gillnet surveys BO- NUS INSPIRE; Wil- lebrand 2015, Öhn- stedt 2015	Salinity, tempera- ture, oxygen, variables from dropvideo: vegetation, sub- strate and occurence of mussels	Baltic Proper, Both- nian Sea, Bothnian Bay	Research project	2014–2015 data from Baltic Proper freely available af- ter BONUS INSPIRE project finishes in 2018	BONUS INSPIRE; Ann-Britt Florin
Cod, sole, eel, lab- rids	Fyke nets	Species distribution modelling	Fredriksson <i>et al.</i> 2010	Salinity, tempera- ture, oxygen, water depth, secchi depth	Kattegat, Skagerrak	Mapping project	Available in Swe- dish reports	Ulf Bergström

Table 19: Sweden (continued)

Species that are mapped and monitored	Observation / Monitoring Method	Mapping Method (brief description)	Reference for methods used	Habitat variables collected	Geographical cov- erage and years of data collection	Purpose	Availability of data or maps (format)	Data owner; contact person
Perch, pikeperch, cyprinids, whitefish etc.	Multimesh gillnets (Nordic coastal)	Data not mapped yet	Survey: "Provfiske i Östersjöns kustom- råden – Djupstrati- fierat provfiske med Nordiska kust- översiktsnät"	Temperature, Sec- chi depth, salinity	Baltic Proper, Both- nian Sea, Bothnian Bay	Several mapping projects + monitor- ing programmes	Data from 2002– 2015 available through SLU data- base KUL	SLU, HaV, NV, Re- gional County Boards; Ulf Bergström
Flounder, turbot	Push nets	Species distribution modelling	Reference for point data: Florin <i>et al.</i> 2009	Meiofauna, macrofauna, epi- fauna, substrate, filamentous algae, turbidity, salinity, structural habitat complexity	Baltic Proper 2006	Research project	No maps	SLU Aqua; Ann-Britt Florin
Habitat mapping targeting vegeta- tion and benthic fauna	Drop video, diving, grab samples	Almost 700 habitat maps of vegetation, benthic fauna, nat- ural values, Helcom underwater bio- topes	See comment	See comment in reference	Baltic Proper, Both- nian Sea, Bothnian Bay	Mapping projects	Free at low resolution	County Boards; AquaBiota Water Research



Ved Stranden 18 DK-1061 Copenhagen K www.norden.org

Essential fish habitats (EFH)

Many fish species in the Baltic Sea are dependent on shallow and sheltered near-shore habitats for their spawning, nursery, feeding and migration. Still, the role of these essential fish habitats, EFH, for the development and support for fish production has received little attention. As coastal EFH often are found in areas heavily impacted by humans, they are subject to many threats and therefore management needs are urgent. EFH also provide and support important ecosystem services and are included in national/international agreements and legislative acts. Despite this, the conservation status of EFH is generally poor in the region. Due to these shortcomings and needs, a workshop was set up to review the importance and protection of as well as threats to coastal EFH in the Baltic Sea. This report describes the outcome of the workshop and future directions for work in this research area.

TemaNord 2016:539 ISBN 978-92-893-4664-1 (PRINT) ISBN 978-92-893-4665-8 (PDF) ISBN 978-92-893-4666-5 (EPUB) ISSN 0908-6692





