

This is an electronic reprint of the original article. This reprint may differ from the original in pagination and typographic detail.

Cell type markers indicate distinct contributions of decidual stromal cells and natural killer cells in preeclampsia

Rytkönen, Kalle T; Adossa, Nigatu; Mahmoudian, Mehrad; Lönnberg, Tapio; Poutanen, Matti; Elo, Laura L

Published in:
Reproduction

DOI:
[10.1530/REP-22-0079](https://doi.org/10.1530/REP-22-0079)

Published: 01/11/2022

Document Version
Final published version

Document License
CC BY

[Link to publication](#)

Please cite the original version:

Rytkönen, K. T., Adossa, N., Mahmoudian, M., Lönnberg, T., Poutanen, M., & Elo, L. L. (2022). Cell type markers indicate distinct contributions of decidual stromal cells and natural killer cells in preeclampsia. *Reproduction*, 164(5), V9-V13. <https://doi.org/10.1530/REP-22-0079>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Cell type markers indicate distinct contributions of decidual stromal cells and natural killer cells in preeclampsia

Kalle T Rytönen^{1,2,*}, Nigatu Adossa^{1,*}, Mehrad Mahmoudian^{1,2}, Tapio Lönnberg^{1,3}, Matti Poutanen² and Laura L Elo^{1,3,4}

¹Turku Bioscience Centre, University of Turku and Åbo Akademi University, Turku, Finland, ²Institute of Biomedicine, Research Centre for Integrative Physiology and Pharmacology, University of Turku, Turku, Finland, ³InFLAMES Research Flagship Center, University of Turku, Turku, Finland and ⁴Institute of Biomedicine, University of Turku, Turku, Finland

Correspondence should be addressed to K T Rytönen; Email: katury@utu.fi

* (K T Rytönen and N Adossa contributed equally to this work)

Abstract

In brief: Preeclampsia is a common serious disorder that can occur during pregnancy. This study uses integrative analysis of preeclampsia transcriptomes and single-cell transcriptomes to predict cell type-specific contributions to preeclampsia.

Abstract: Preeclampsia is a devastating pregnancy disorder and a major cause of maternal and perinatal mortality. By combining previous transcriptomic results on preeclampsia with single-cell sequencing data, we here predict distinct and partly unanticipated contributions of decidual stromal cells and uterine natural killer cells in early- and late-onset preeclampsia.

Reproduction (2022) **164** V9–V13

Preeclampsia is commonly classified as early-onset preeclampsia (EOP) (before week 34) and late-onset preeclampsia (LOP). EOP is characterized by defective placentation and spiral remodeling, whereas in LOP, normal placentation has apparently taken place, but placental capacity is exceeded and/or other factors contribute to the symptoms. Recent studies suggest that maternal cell types, such as endometrial stromal cells, have a more critical role in the etiology of preeclampsia than previously anticipated (Conrad *et al.* 2017, Garrido-Gomez *et al.* 2017). However, the current models of preeclampsia only loosely explain their relative contributions and the cell type-specific molecular mechanisms.

Here, we conduct an analysis that combines decidual bulk transcriptomic data from preeclampsia with recently published single-cell sequencing (scRNA-seq) data from healthy women. Specifically, we utilized the previously detected differentially expressed genes for severe EOP and severe LOP from decidua samples from term pregnancies (Tong *et al.* 2018), and from secretory phase of the menstrual cycle of women that previously experienced severe preeclampsia (Garrido-Gomez *et al.* 2021). These were intersected with the cell type marker lists from scRNA-seq transcriptomic atlases from endometrium taken during the secretory phase of menstrual cycle (Wang *et al.* 2020), decidua from women undergoing selective pregnancy termination

during the first trimester (Vento-Tormo *et al.* 2018), and decidua from women at term pregnancy (Pique-Regi *et al.* 2019). With this integrative analysis, we were able to predict distinct cell type-specific contributions to the pathogenesis of both EOP and LOP.

We first re-ran the cluster analysis for the three scRNA-seq datasets using Seurat 4.0 (Hao *et al.* 2021), re-annotated the main uterine cell types concordantly (Fig. 1A, B and C), identified cell type-specific marker genes and combined these from the three studies to an extended cell type marker list. To search for the presence of cell type-specific markers among the preeclampsia-regulated genes, we intersected scRNA-seq marker lists and disease genesets and conducted Fisher's exact tests (Fig. 1D). We discovered two robust overrepresentation signals: LOP downregulated genes were enriched with endometrial/decidual stromal cell (dS) markers (combined uterine markers $P=1.5 \times 10^{-32}$, 5.1-fold) and EOP upregulated genes were enriched with uterine natural killer cell (uNK) markers ($P=1.3 \times 10^{-22}$, 8.6-fold) (Fig. 1D). This suggests that LOP and EOP have distinct cell type-specific etiological characteristics. While LOP is associated with transcriptomic changes in stromal cells, EOP is more closely associated with transcriptomic changes in uNKs. As decidualization takes place spontaneously during the non-pregnant menstrual cycle, these results also suggest that the defects in uterine differentiation that contribute to preeclampsia may be




Figure 1 Preeclampsia regulated genes are enriched with decidua stromal cell and uterine natural killer cell marker genes. The uterine cell types in (A) secretory phase menstrual cycle (Wang *et al.* 2020), (B) first trimester pregnancy (Vento-Tormo *et al.* 2018) and (C) term not in labor pregnancy (Pique-Regi *et al.* 2019) single-cell transcriptomes with harmonized annotation. For (A) and (B), the preprocessed scRNA-seq UMI counts were extracted from ArrayExpress accession E-MTAB-6701 (Vento-Tormo *et al.* 2018), including 6 decidua samples (6–12 weeks of

Figure 1 Continued

gestation) with a total of 36,186 cells, and GEO accession GSE111976 (Wang *et al.* 2020), including 10 endometrial samples (cycle days 16–26) with a total of 71,032 cells, respectively. For (C), the raw data were extracted from dbGaP accession phs001886.v1.p1 (Pique-Regi *et al.* 2019) with consent, only term no labor (basal plate, chorioamniotic membranes) samples were selected with a total of 13,730 cells. All three datasets were produced using 10× Genomics scRNA-seq. Samples were pre-processed using cellranger-3.1.0's count function with default parameters and prebuilt reference genome (hg38). We used the standard analysis protocol of Seurat 4.0 (Hao *et al.* 2021) and the annotation of the clusters was harmonized to follow the one from the first trimester study (Vento-Tormo *et al.* 2018). (D) A heatmap of Fisher's exact test *P* values for the overlaps of uterine cell type marker genes and preeclampsia differentially expressed genes. The arrows mark the subsets that are up- or downregulated genes in the original studies. The marker gene lists used for each main cell type were combined lists from the condition-specific cell type markers from A, B and C. Early-onset (before week 34) severe preeclampsia (EOP) (*n* = 3), late-onset severe preeclampsia (LOP) (*n* = 3) and control (*n* = 3) samples were from term deciduas delivered by cesarean section (Tong *et al.* 2018). Samples for women with previous severe preeclampsia (M-PP) (*n* = 17) together with controls (*n* = 12) were from late secretory menstrual cycle (days 22–32) endometrium (Garrido-Gomez *et al.* 2021). Statistical significance of the enrichment was determined using Fisher's exact test in R, with all Refseq protein-coding genes (*n* = 20,203) as the background. (E) A heatmap of Fisher's exact test *P* values for the overlaps of the preeclampsia differentially expressed genes and the subpopulations of endometrial/decidual stromal cells (dS1, dS2 and dS3) from the first trimester study. (F) Genes in the LOP downregulated – stromal dS3 marker intersection (first trimester). (G) Trajectory inference of differentiating dS cells using Slingshot (Street *et al.* 2018). Slingshot uses minimum spanning tree algorithm to reconstruct the lineage structure among differentiating cells from scRNA-seq data. In the 2D visualization, the axes represent the first principal components. The black dots represent the cells in the lower dimension, and the arrow-headed lines depict the suggested differentiation trajectories estimated by Slingshot. (H) Decorin (*DCN*) and (I) galectin 1 (*LGALS1*) expression across the dS cell trajectories. (J) Genes in the EOP upregulated – uNK cell type marker intersection. (K) Trajectory inference of uNK cell subpopulations from the first trimester study using Slingshot. (L) *CD7* expression across the uNK trajectories. For (E–L) only the first trimester data was used. For (F) and (J), the top functional enrichment categories (GO, Hallmark) are displayed on the left. Functional enrichment analysis was done using the Metascape tool with all available pathway databases (GO biological processes, Reactome Gene Sets, Canonical Pathways, Biocarta Gene Sets, WikiPathways, KEGG Pathways and Hallmark Gene Sets). Trajectory inference was done using the `infer_trajectory` function of Slingshot in the 'dyno' version '0.1.2' (Saelens *et al.* 2019).

detectable already in the non-pregnant menstrual cycle. However, in the analyzed dataset of genes differentially expressed in the secretory phase of the menstrual cycle from women that previously had preeclampsia (Garrido-Gomez *et al.* 2021), the most significant signature was observed for perivascular cells with less striking enrichment ($P = 2.0 \times 10^{-10}$, 3.7-fold) compared to those we observed with EOP and LOP (Tong *et al.* 2018).

We next focused on the cell subpopulations among stromal cells in the first trimester data (Vento-Tormo *et al.* 2018). The three annotated subpopulations (dS1, dS2 and dS3) of stromal cells reflect the stages of the decidualization (differentiation), and dS3 represents decidualized cells with high prolactin expression (Vento-Tormo *et al.* 2018). We repeated the Fisher's exact tests separately for dS1, dS2 and dS3 (Fig. 1E) and observed enrichment of dS3 markers among the LOP downregulated genes ($P = 3.7 \times 10^{-38}$, 16-fold). This suggests that in LOP the normal decidualization of maternal stromal cells is defected, further supporting the recent observation by others on severe preeclampsia (Rabaglino *et al.* 2015, Garrido-Gomez *et al.* 2017, 2021). These dS3 – LOP downregulated genes had progressively higher expression during decidualization in healthy donors (Fig. 1F), and they were functionally enriched with terms such as 'epithelial–mesenchymal transition' (EMT) (Hallmark, $P = 6.1 \times 10^{-8}$) and 'wounding' (GO, $P = 5.9 \times 10^{-5}$). To visualize the decidualization trajectory and the expression of selected genes associated with these terms in healthy donors, we used Slingshot (Street *et al.* 2018) (Fig. 1G, H and I). For instance, the expression levels of decorin (*DCN*) and galectin 1 (*LGALS1*) are markedly increased during the decidualization and have the

highest expression in dS3 (Fig. 1H and I). *DCN* is a decidualization marker and *LGALS1* enhances maternal immunotolerance during pregnancy. On the other hand, LOP upregulated genes were overrepresented with dS1 marker genes ($P = 1.8 \times 10^{-11}$, 9.9-fold) (Fig. 1E), including fibroblast marker, actin alpha 2 (*ACTA2*), for the non-decidualized stage, being again in line with the prediction that decidualization defects contribute to LOP. We also observed that the EOP downregulated genes were similarly enriched with dS3 markers ($P = 1.3 \times 10^{-11}$, 7.6-fold), suggesting that gene regulatory changes associated with decidualization defects of stromal cells also contribute to EOP, but not to the same extent as in LOP.

For uNK subpopulations (dNK1, dNK2 and dNK3) (Vento-Tormo *et al.* 2018), we did not detect robust subpopulation-specific overrepresentation among EOP or LOP regulated genes. We hypothesized that this is because the subpopulation clusters are only moderately separated from each other. Thus, we conducted a functional enrichment analysis of all the uNK cell type marker genes that were upregulated in EOP (Fig. 1D) and found terms such as 'Allograft rejection' (Hallmark, $P = 4.9 \times 10^{-8}$) and 'Leukocyte activation' (GO, $P = 5.1 \times 10^{-8}$) (Fig. 1J), which are generally associated with reduced maternal immunotolerance. Although in healthy donors these uNK marker genes did not present as clear differentiation trajectory associated expression patterns as observed in stromal cells, we observed a modest trend of genes in 'Allograft rejection' to be more expressed in the dNK2 and dNK3 populations compared to uNK1 (Fig. 1J), including, e.g. *CD7* (Fig. 1K and L). This suggests that uNK markers upregulated in

the more differentiated subpopulations (dNK2 and dNK3) may associate with reduced immunotolerance and preeclampsia.

Finally, for the top overrepresentation gene intersect for women with previous preeclampsia (Garrido-Gomez *et al.* 2021), the perivascular cells – preeclampsia downregulated signature, we also conducted a functional enrichment analysis. The detected top terms were ‘EMT’ (Hallmark, $P=4.4 \times 10^{-9}$) and ‘blood vessel development’ (GO, $P=7.8 \times 10^{-9}$). Notably, EMT term was also detected for the stromal dS3 – LOP downregulated intersect, and the perivascular cell clusters have been associated with endometrial stromal stem cells (Queckbörner *et al.* 2021) that have previously been reported to underlie reproductive disorders.

The main limitations of this study involve the temporal specificity in the dataset intersections. First, EOP and LOP data (Tong *et al.* 2018) were collected upon delivery and the transcriptomic signatures upon delivery may be different from the initial signatures for EOP and LOP earlier during the pregnancy (Rabaglino & Conrad 2019). Second, we selected to combine the scRNA-seq cell type-specific markers from the three different time points in order to have more comprehensive cell type marker lists, but this inevitably decreased the temporal specificity of these lists for a given one time point.

In conclusion, our analysis with single-cell data-derived marker genes extends the resolution of previous heterogeneous EOP vs LOP comparisons. We observed that preeclampsia downregulated genes (especially in LOP) are enriched with the markers of decidualized stromal cells (dS3) indicating a critical role for decidualization defects in the etiology of preeclampsia. Our results support previous observations on the importance of decidualization defects (Rabaglino *et al.* 2015, Garrido-Gomez *et al.* 2017, 2021). Notably, these studies collected severe preeclampsia samples without classification to EOP and LOP, whereas our analysis presents a novel observation that the overrepresentation signal for stromal cell decidualization defects is stronger in LOP rather than EOP. Curiously, decidual stromal cells have been suggested to have independently evolved two of their main functions; first, the anti-inflammatory reaction associated with implantation in the stem of placental mammals, and secondly, the ability to support extended maintenance of pregnancy in Euarchontoglires (suprprimates) including humans (Chavan *et al.* 2016). The robust LOP-specific signal may be associated with the function of decidual stromal cells to support the maintenance of pregnancy rather than the initial anti-inflammatory role in implantation and early placentation. These results, therefore, emphasize the abovementioned dual role of stromal cells in reproductive health. On the other hand, uNK markers linked with reduced immunotolerance were specifically enriched in the EOP upregulated genes which is in line with the critical role of uNK cells to promote immunotolerance during spiral

artery modeling and placentation. Overall, our analysis highlights the potential of and motivates future single-cell studies on preeclampsia.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this article.

Funding

This work was supported by Päivikki and Sakari Sohlberg Foundation (K T R), Emil Aaltonen Foundation (K T R), University of Turku Graduate School (N A), Sigrid Juselius Foundation (M P and L L E) and Academy of Finland (L L E).

Author contribution statement

K T R designed the work. K T R, N A and M M performed analysis. K T R, N A, M M, T L, M P and L L E wrote the paper.

References

- Chavan AR, Bhullar BA & Wagner GP 2016 What was the ancestral function of decidual stromal cells? A model for the evolution of eutherian pregnancy. *Placenta* **40** 40–51. (<https://doi.org/10.1016/j.placenta.2016.02.012>)
- Conrad KP, Rabaglino MB & Post Uiterweer ED 2017 Emerging role for dysregulated decidualization in the genesis of preeclampsia. *Placenta* **60** 119–129. (<https://doi.org/10.1016/j.placenta.2017.06.005>)
- Garrido-Gomez T, Dominguez F, Quiñonero A, Diaz-Gimeno P, Kapidzic M, Gormley M, Ona K, Padilla-Iserte P, McMaster M, Genbacev O *et al.* 2017 Defective decidualization during and after severe preeclampsia reveals a possible maternal contribution to the etiology. *PNAS* **114** E8468–E8477. (<https://doi.org/10.1073/pnas.1706546114>)
- Garrido-Gomez T, Castillo-Marco N, Clemente-Ciscar M, Cordero T, Muñoz-Blat I, Amadoz A, Jimenez-Almazan J, Monfort-Ortiz R, Climent R, Perales-Marin A *et al.* 2021 Disrupted PGR-B and ESR1 signaling underlies defective decidualization linked to severe preeclampsia. *eLife* **10** e70753. (<https://doi.org/10.7554/eLife.70753>)
- Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, Lee MJ, Wilk AJ, Darby C, Zager M *et al.* 2021 Integrated analysis of multimodal single-cell data. *Cell* **184** 3573.e29–3587.e29. (<https://doi.org/10.1016/j.cell.2021.04.048>)
- Pique-Regi R, Romero R, Tarca AL, Sandler ED, Xu Y, Garcia-Flores V, Leng Y, Luca F, Hassan SS & Gomez-Lopez N 2019 Single cell transcriptional signatures of the human placenta in term and preterm parturition. *eLife* **8** e52004. (<https://doi.org/10.7554/eLife.52004>)
- Queckbörner S, von Grothhusen C, Boggavarapu NR, Francis RM, Davies LC & Gemzell-Danielsson K 2021 Stromal heterogeneity in the human proliferative endometrium – a single-cell RNA sequencing study. *Journal of Personalized Medicine* **11** 448. (<https://doi.org/10.3390/jpm11060448>)
- Rabaglino MB & Conrad KP 2019 Evidence for shared molecular pathways of dysregulated decidualization in preeclampsia and endometrial disorders revealed by microarray data integration. *FASEB Journal* **33** 11682–11695. (<https://doi.org/10.1096/fj.201900662R>)
- Rabaglino MB, Uiterweer EDP, Jeyabalan A, Hogge WA & Conrad KP 2015 Bioinformatics approach reveals evidence for impaired endometrial maturation before and during early pregnancy in women who developed preeclampsia. *Hypertension* **65** 421–429. (<https://doi.org/10.1161/HYPERTENSIONAHA.114.04481>)
- Saelens W, Cannoodt R, Todorov H & Saeys Y 2019 A comparison of single-cell trajectory inference methods. *Nature Biotechnology* **37** 547–554. (<https://doi.org/10.1038/s41587-019-0071-9>)

- Street K, Riso D, Fletcher RB, Das D, Ngai J, Yosef N, Purdom E & Dudoit S** 2018 Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC Genomics* **19** 477. (<https://doi.org/10.1186/s12864-018-4772-0>)
- Tong J, Zhao W, Lv H, Li WP, Chen ZJ & Zhang C** 2018 Transcriptomic profiling in human decidua of severe preeclampsia detected by RNA sequencing. *Journal of Cellular Biochemistry* **119** 607–615. (<https://doi.org/10.1002/jcb.26221>)
- Vento-Tormo R, Efremova M, Botting RA, Turco MY, Vento-Tormo M, Meyer KB, Park JE, Stephenson E, Polański K, Goncalves A *et al.*** 2018 Single-cell reconstruction of the early maternal–fetal interface in humans. *Nature* **563** 347–353. (<https://doi.org/10.1038/s41586-018-0698-6>)
- Wang W, Vilella F, Alama P, Moreno I, Mignardi M, Isakova A, Pan W, Simon C & Quake SR** 2020 Single-cell transcriptomic atlas of the human endometrium during the menstrual cycle. *Nature Medicine* **26** 1644–1653. (<https://doi.org/10.1038/s41591-020-1040-z>)

Received 4 March 2022

First decision 28 April 2022

Revised manuscript received 18 August 2022

Accepted 15 September 2022